



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, C07K 14/705, 16/28, G01N 33/50, C12N 15/62, G01N 33/53, C12Q 1/68</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/32627</b> <b>(43) International Publication Date:</b> 1 July 1999 (01.07.99)
<b>(21) International Application Number:</b> PCT/US98/27400 <b>(22) International Filing Date:</b> 23 December 1998 (23.12.98) <b>(30) Priority Data:</b> 60/068,690 23 December 1997 (23.12.97) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 60/068,690 (CIP) Filed on 23 December 1997 (23.12.97)		<b>(74) Agents:</b> CARROLL, Peter, G. et al.; Medlen & Carroll, LLP, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US). <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
<b>(71) Applicant (for all designated States except US):</b> THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SAMELSON, Lawrence, E. [US/US]; 6707 East Avenue, Chevy Chase, MD 20815 (US). ZHANG, Weiguo [-/US]; 259 Congressional Lane #410, Rockville, MD 20852 (US).		<b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	

(54) Title: THE PROTEIN TYROSINE KINASE SUBSTRATE LAT AND ITS USE IN THE IDENTIFICATION OF (ANT)AGONISTS OF THE KINASE

```
human LAT      1 MEZAILVPCVLGLLLLPIL.AMLHALCVHCHRLPGSDSTSSDSLPRGI 49  
               1 ... | .|. |. |. |. |. |. |. |. |. |. |. |. |. |  
murine LAT     1 HEADALSPVGLGLLLLPFLVTLLAALCVRCRELPSDSTSTESLPRSI 50  
  
50 QFKRPHTVAPWPPA...PFVTSYPPLSQPDLLPIRSPQFLOGSERTPSS 97  
   :.|.:...|.:|| | |:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|  
51 LIKPPQITVPRTPAVSPLVTSFPPLRQPDLLPIRSPQFLOGSHMPSS 100  
  
98 RRDSDGANSVASINEEPACK..DADEDEDENHPGLVVLPDSTPATST 145  
  ..|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|  
101 QQNSDDANSVASINQEPAKNVDADADEDDPN.GLVVLPDSSPAAVP 149  
  
146 AAPSA PALSTPGIRDSAFSMESIDDVNVPESGESAEASLDGSREYNVS 195  
  ...|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|  
150 VVSSAPVPSNPDLDGSAFSVESCEDVNVPSESESAAEASLDGSREYNVS 199  
  
196 QELHPGAAKTEPAALSSQEAEE.....VEKEQAPDENLQELN 233  
    -|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|:|  
200 PEQQP.VTRAELASVNSQEVEDEGEREGVDGEKAPDENLQELN 242
```

**Human (1-233) and Murine LAT Amino Acid Sequence (1-242)**

**(57) Abstract**

The invention generally relates to compositions and methods for identifying and testing tyrosine kinase signaling pathway agonists and antagonists, and more particularly, methods and compositions for screening compounds and identifying compounds that will modulate the interaction of protein tyrosine kinase substrates with their intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CJ	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THE PROTEIN TYROSINE KINASE SUBSTRATE LAT AND ITS USE IN THE IDENTIFICATION OF (ANT)AGONISTS OF THE KINASE

5           This invention was made with government support from a grant from the National Institute of Child Health and Human Development (NICHD). The United States Government has certain rights in the invention.

**FIELD OF THE INVENTION**

10           The invention generally relates to compositions and methods for identifying and testing tyrosine kinase signaling pathway agonists and antagonists, and more particularly, methods and compositions for screening compounds and identifying compounds that will modulate the interaction of protein tyrosine kinase substrates with their intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway.

15           **BACKGROUND OF INVENTION**

          The immune system has evolved to provide an organism with a flexible and dynamic mechanism to respond specifically to a wide variety of antigenic insults. In order for an immune response to occur following antigenic challenge, the antigen must not only be  
20           recognized by antigen specific lymphocytes, but this recognition event must lead to a variety of cellular responses. T lymphocytes, together with B lymphocytes, represent the two antigen specific components of the cellular immune system. T cells are central to the immune response and control the response to infectious agents, tumors and transplants.

          The T cell receptor for antigen (TCR) recognizes and binds foreign antigens. This  
25           binding event leads to T cell activation. The activation of T lymphocytes is a complex process which results in cell growth and differentiation. The engagement of the TCR on mature peripheral T cells initiates multiple intracellular signals that can lead to cellular proliferation and the acquisition of complex effector functions.

          The biochemical mechanisms that couple receptor binding to these intracellular events  
30           have been intensively investigated. Early events such as activation of tyrosine phosphorylation, elevation of intracellular calcium, activation of lipid-dependent kinases, and activation of Ras and its downstream kinase cascade are well known [Weiss, A. and Littman, D. R., Signal transduction by lymphocyte antigen receptors, *Cell* 76:263-274 (1994); Cantrell,

D., T Cell Antigen Receptor Signal Transduction Pathways, *Annu. Rev. Immunol.* 14:259-274 (1996)]. Moreover, there has been analysis of the events involved in transcriptional induction of T cell-specific lymphokines. There remain, however, significant gaps in the understanding of TCR signaling, particularly in how the early tyrosine phosphorylation events couple  
5 receptor activation to later cellular events.

Perhaps the most critical insight of the past decade in the study of signal transduction has been the recognition that activation of receptor tyrosine kinases results in the assembly of multimolecular complexes at the cytoplasmic domain of the receptor. Likewise, a major role of the tyrosine phosphorylation cascade initiated by TCR engagement is the assembly of  
10 multimolecular signaling complexes at and near the TCR itself. The identity and functions of these proteins that engage in such molecular interactions, are critical in the understanding of T cell receptor signal transduction.

## SUMMARY OF THE INVENTION

15 The invention generally relates to compositions and methods for identifying and testing tyrosine kinase signaling pathway agonists and antagonists, and more particularly, methods and compositions for screening compounds and identifying compounds that will modulate the interaction of protein tyrosine kinase substrates with their intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway. It  
20 is not intended that the present invention be limited to particular signaling pathways. The present invention contemplates that the methods and compositions described herein will be useful for identifying protein tyrosine kinase (PTK) downstream signaling proteins, particularly in cells including but not limited to T cells, NK cells, and mast cells, and will enable the identification of compounds that will modulate the interactions of protein tyrosine  
25 kinase that bind particular intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway.

In preferred embodiments, "Linker for Activation of T-cells" (LAT, and in particular, fragments of LAT) are useful in drug screening assays designed to identify drugs that interfere with the specific binding of Grb2 or GST-GRb2 with LAT and thereby block  
30 the activation of downstream signaling molecules.

In one embodiment, the present invention contemplates screening compounds and identifying compounds that modulate the interactions of T cell protein kinases, and their substrates, in particular ZAP-70 and/or Syk tyrosine kinase.

Furthermore, the present invention contemplates identifying ZAP-70 and/or Syk substrates and ZAP-70 and/or Syk-binding ligands, and compounds that will modulate the interaction of ZAP-70 and/or Syk kinase with ZAP-70 and/or Syk substrates and ZAP-70 and/or Syk-binding ligands.

5 In one embodiment, the present invention contemplates identifying compounds that modulate the interaction of LAT, which binds to activated, Tyr phosphorylated ZAP-70 and/or Syk kinases.

In preferred embodiments, LAT (and, in addition, fragments of LAT and mutated LAT) are useful in drug screening assays designed to identify drugs that interfere with the specific binding activated ZAP-70 and/or Syk kinases with their substrates and thereby block the activation of downstream signaling molecules.

10 In other embodiments, the present invention contemplates identifying compounds that modulate the interaction of LAT, which may bind to kinases other than ZAP-70 and/or Syk. In other embodiments, the invention provides an isolated LAT polypeptide, or a fragment thereof, having ZAP-70 and/or Syk kinase-specific binding affinity. The invention provides nucleic acids encoding the LAT polypeptide and LAT fragments as part of expression vectors for introduction into cells. The invention provides methods of identifying intracellular molecules which interact with LAT or LAT fragments, as well as exogenous agents (i.e. drugs) which disrupt or enhance the binding of LAT and/or fragments thereof to such intracellular targets.

20 The claimed polypeptide LAT finds particular use in screening assays for agents or lead compounds for agents useful in the diagnosis, prognosis or treatment of disease, particularly disease associated with undesirable cell growth, differentiation, proliferation and T cell anergy. One such assay involves forming mixtures of 1) LAT (or fragments thereof) and 2) an intracellular LAT-binding ligand, in the presence or absence of 3) a prospective drug candidate. The mixtures are made under conditions that permit the binding of the intracellular LAT-binding ligand to LAT (or fragments thereof) and the mixtures are then analyzed for the presence of such binding. A difference in such binding in the presence of such a drug candidate indicates that the agent is capable of modulating the binding of LAT (or fragments thereof) to an intracellular LAT-binding ligand.

30 It is not intended that the present invention be limited by the species (human, murine, rat, etc.) of the binding ligands described above. The polypeptide LAT and LAT fragments may bind across species. Moreover, the nucleic acid sequences described herein allow for the

identification of homologues in other species by various methods, including but not limited to amplification (e.g. PCR) using primers designed from the nucleic acid sequence of one species (e.g. mouse) on the nucleic acid template of another species (e.g. human).

In one embodiment, the present invention contemplates an isolated nucleic acid encoding at least a fragment of a protein having the amino acid sequence set forth in SEQ ID NO:4. It is not intended that the present invention be limited by the size or nature of the fragment (although it is preferred that such fragments are capable of binding kinases). In one embodiment, said nucleic acid comprises SEQ ID NO:1 and encodes full-length LAT as set forth in (SEQ ID NO:4). In yet another embodiment, said nucleic acid encodes a fusion protein.

It is not intended that the present invention be limited as to the specific nature of the nucleic acid encoding the peptides described above. In one embodiment, said nucleic acid is contained in a vector. In another embodiment, said vector is in a host cell.

The present invention also contemplates complexes of ligands. In one embodiment, the present invention contemplates a composition, comprising a LAT -kinase complex comprising a purified peptide having at least a portion of the amino acid sequence set forth in SEQ ID NO:4 specifically bound to activated ZAP-70 and/or Syk (or other kinases). Again, the peptides bound specifically to activated ZAP-70 and/or Syk kinases may be full-length LAT or a fragment defined by a portion of the amino acid sequence as set forth in SEQ ID NO:4. The peptide may be part of a fusion protein. The complex can also contain other ligands, such as effector molecules downstream to PTKs. The complexes can be used to identify other ligands (as described below).

As noted above, the present invention contemplates compound screening assays. In one embodiment, the present invention contemplates a method for compound screening, comprising: a) providing: i) a peptide comprising at least a portion of the amino acid sequence set forth in SEQ ID NO:4, wherein said portion is capable of binding to ZAP-70 and/or Syk kinases, ii) ZAP-70 and/or Syk kinases, and iii) one or more compounds for screening; b) mixing, in any order, said peptide, said ZAP-70 and/or Syk kinases and said one or more compound; and c) measuring the extent of binding of said peptide to said ZAP-70 and/or Syk kinases. Again, the peptides may be full-length LAT or mutated LAT or a fragment defined by a portion of the amino acid sequence as set forth in SEQ ID NO:4. The peptide may also be part of a fusion protein. The present invention also contemplates embodiments where either the peptide or kinase is bound to other ligands, such as effector

molecules downstream to PTKs or other PTKs. These complexes can be used in the compound screening assay described above.

The present invention specifically contemplates an isolated polypeptide comprising at least a portion of the amino acid sequence of SEQ ID NO:4. It is not intended that the present invention be limited to a specific portion. However, in one embodiment, said portion comprises a region comprising at least one tyrosine. In another embodiment, said portion comprises more than one tyrosine (and more preferably more than five tyrosine residues). The present invention also contemplates domains of the LAT protein. While not limited to any particular domain, domains such as the cytosolic domain (defined approximately by amino acids 28 to 233).

The present invention also specifically contemplates antibodies (both polyclonal and monoclonal) to LAT, LAT fragments, mutant LAT, mutant LAT fragments, and LAT/LAT Binding Ligand complexes. In one embodiment, the present invention contemplates A purified antibody which binds specifically to a polypeptide comprising at least a portion of the amino acid sequence of SEQ ID NO:4.

The present invention also contemplates an isolated polynucleotide encoding the polypeptide comprising the sequence of SEQ ID NO:4. In one embodiment, said polynucleotide comprises the sequence of SEQ ID NO: 1. The present invention contemplates embodiments where said polynucleotide is contained on a recombinant expression vector and where said expression vector containing said polynucleotide sequence is contained within a host cell.

In some embodiments, the present invention contemplates nucleic acids capable of hybridizing to portions of the LAT gene. In one embodiment, the present invention contemplates a polynucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO:1.

As discussed above, screening methods are contemplated. In a preferred embodiment, the present invention contemplates a method of screening a compound, said method comprising: a) providing, in any order: i) a peptide comprising at least a portion of the amino acid sequence set forth in SEQ ID NO:4, wherein said portion is capable of binding to a LAT binding ligand; ii) a LAT binding ligand; and iii) one or more compounds for screening; b) mixing, in any order, said peptide, said LAT binding ligand and said one or more compound; and c) measuring the extent of binding of said peptide to said LAT binding ligand.

It is not intended that the present invention be limited to a particular LAT binding ligand. In one embodiment, said LAT binding ligand comprises a tyrosine kinase. In another embodiment, said kinase comprises ZAP-70 kinase. In yet another embodiment, said kinase comprises Syk kinase.

In the embodiments described above, it may be useful to use the entire LAT polypeptide or fragments. In addition, it may be useful where said peptide is part of a fusion protein.

The present invention also contemplates detecting LAT (e.g. detecting LAT in cells and other biological samples). In one embodiment, the present invention contemplates a method for detecting the presence of a portion of the polypeptide having the amino acid sequence set forth in SEQ ID NO:4, said method comprising the steps of: a) providing in any order: i) an antibody capable of reacting with a portion of the polypeptide having the sequence set forth in SEQ ID NO:4; and ii) a sample suspected of containing at least a portion of the polypeptide having the sequence set forth in SEQ ID NO: 4; b) combining said antibody and said sample under conditions such that a complex is formed between said antibody and said portion of said polynucleotide; and c) detecting said complex. Again, the antibody can be polyclonal antibody or monoclonal antibody. A variety of cells and samples are contemplated, including but not limited to NK cells, mast cells and lymphocytes.

The present invention also contemplates using antibodies in other formats. In one embodiment, LAT is detected on gels by immunoblotting.

The present invention is not limited to antibody detection. It may be useful to detect nucleic acid encoding the LAT polypeptide or detect gene expression. In one embodiment, the present invention contemplates a method for detecting the presence of polynucleotide sequences encoding at least a portion of LAT gene in a sample, said method comprising the steps of: a) providing in any order: i) a polynucleotide comprising a sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO:1; and ii) a sample suspected of containing nucleic acid comprising the sequence of SEQ ID NO:1; b) combining said polynucleotide and said sample under conditions such that a hybridization complex is formed between said polynucleotide and said sample nucleic acid; and c) detecting said hybridization complex. The present invention contemplates detecting both RNA and DNA. Again, a variety of cells and samples can be used, including but not limited to, NK cells, mast cells and lymphocytes. It is desirable that said detected hybridization complex correlates with expression of the LAT gene in said lymphocytes.



## DESCRIPTION OF FIGURES

FIGURE 1 shows the purification, cDNA cloning, and the deduced amino acid sequence of LAT (Linker for activation of T cells)

FIG. 1(A) is a representative immunoblot of Jurkat cells stimulated with C305 (+) or unstimulated cells (-), analyzed by immunoblotting with anti-phosphotyrosine antibody (PY20).

FIG. 1(B) shows the resolution by reverse-phase microbore HPLC of the affinity-purified p36-38 protein (isolated from stimulated Jurkat cells). Five peptide peaks (fraction 31, 40, 48, 55, 82) were selected for mass spectrometry and microsequencing.

FIG. 1(C) shows the analysis of peptides from the indicated peaks. The single amino acid code is used and with the following additional: X=no assignment, uppercase=full confidence assignment, lowercase=ambiguous residue and n.d.=not determined.

FIGURE 2 shows the characterization of the LAT protein.

FIG. 2(A) is a representative immunoblot of 293T cells transfected with empty vector (pcDNA3) or with myc-tagged LAT in pcDNA3 (+). Post-nuclear lysates of 293T cells were immunoprecipitated with anti-myc antibody (9E10), resolved on SDS-PAGE and immunoblotted with anti-myc antibody.

FIG. 2(B) is a representative immunoblot of lysates from C305 stimulated Jurkat T cells. Proteins immunoprecipitated with rabbit anti-LAT antibody, were treated with calf intestine alkaline phosphatase where indicated, and blotted with anti-phosphotyrosine (pY) and anti-LAT antibodies.

FIG. 2(C) depicts the kinetics of LAT phosphorylation in OKT3 stimulated Jurkat cells. Lysates were immunoprecipitated with anti-LAT antibody, immunoprecipitated proteins were resolved on SDS-PAGE, and blotted with anti-phosphotyrosine (pY) and anti-LAT antibodies.

FIG. 3. shows the tissue distribution of LAT.

FIG. 3(A) is a representative Northern blot analysis of poly (A)<sup>+</sup> RNA from different human tissues.

FIG. 3(B) depicts the Northern blot analysis of total RNA from different cell lines: Jurkat (T cell), YT (NK-like cell), THP1 (monocyte), Raji (B lymphoma), Jiyoye (B lymphoma), K562 (myelomonocytic cells), HeLa (fibroblastoid) and RBL (mast cell). The same membrane was also hybridized with a  $\beta$ -actin probe.

FIGURE 4 shows LAT is phosphorylated by PTKs and associates with Grb2, Grap, the p85 subunit of PI-3K and PLC- $\gamma$ 1 in 293T cells.

FIG. 4(A) is a representative immunoblot showing tyrosine phosphorylation of LAT by Syk and ZAP-70 PTKs. FLAG-tagged LAT in pcDNA3 was transfected in 293T cells either alone or cotransfected with other PTKs as indicated.

FIG. 4(B) is a representative immunoblot, showing association of LAT with signaling molecules in 293T cells. FLAG-tagged LAT was cotransfected with pCEFL/HA-Grb2, pSR $\alpha$ /Grap-myc, pcDNA3/HA-p85 and pcDNA3/PLC- $\gamma$ 1 in the absence or presence of ZAP-70 and Lck PTKs. Brij lysates of transfected 293T cells were immunoprecipitated with anti-FLAG, and association of LAT with tested proteins was determined by blotting with specific antibodies.

FIG. 4(C) is a representative immunoblot, showing LAT associates with Grb2 or Grap by reciprocal immunoprecipitation. Lck, ZAP-70, and LAT were cotransfected with either Grb2-HA or Grap-myc as indicated, and 293T cell lysates were immunoprecipitated with anti-FLAG and anti-HA antibodies if cotransfected with Grb2-HA, or anti-FLAG and anti-Myc antibodies if cotransfected with Grap-myc.

FIGURE 5 shows LAT associates with Grb2, p85 and PLC- $\gamma$ 1 in Jurkat T cells.

FIG. 5(A) is a representative immunoblot, showing GST-Grb2 associates with LAT through its SH2 domain. GST fusion proteins cross-linked to glutathione-agarose were used to precipitate proteins from unstimulated or stimulated with C305 Jurkat lysates. GST-Grb2 (SH2\*) has a mutation in the SH2 domain (R86K). GST-Grb2 (SH3\*) has mutations at both N-terminal and C-terminal SH3 domains (P49L/G203R).

FIG. 5(B) is a representative immunoblot, depicting lysates from unstimulated or C305 stimulated Jurkat T cells subjected to anti-Grb2 or anti-PLC- $\gamma$ 1 immunoprecipitation, and immunoblotted with anti-LAT antibody.

FIG. 5(C) is a representative immunoblot, depicting lysates from Jurkat T cells subjected to preimmune serum or anti-LAT antibody immunoprecipitation, resolved on SDS-PAGE and immunoblotted with anti-phosphotyrosine antibody (4G10). The same membrane was stripped and also blotted with anti-Grb2, p85, Vav, Cbl, PLC- $\gamma$ 1 to identify specific proteins. Only anti-p85 blotting is shown in this figure.

FIGURE 6 shows overexpression of a mutant form of LAT (Y171/I91F) blocks complex formation and TCR mediated AP-1 and NFAT transcriptional activation.

FIG. 6(A) Jurkat cells were stably transfected with wild-type (WT) or mutant (Y171/191F) LAT. Exogenous LAT was immunoprecipitated with anti-myc antibody from cells either left unstimulated or stimulated with C305 (1:50). Samples were analyzed on SDS-PAGE and blotted with 4G10. The same membrane was also blotted with anti-PLC- $\gamma$ 1, anti-p85, anti-Grb2, anti-SOS, and anti-LAT after stripping.

FIG. 6(B) shows secreted alkaline phosphatase (SEAP) activity and the data shown is representative of three independent experiments. Jurkat TAG cells were cotransfected with 20  $\mu$ g pSX-AP-1/SEAP with 20  $\mu$ g empty vector, pEF/LAT(WT), or pEF/LAT(Y171/191F), and then stimulated with OKT3 (1:1000), or stimulated with PMA and ionomycin, or left unstimulated.

FIG. 6(C) shows SEAP activity and the data shown is representative of three independent experiments. Jurkat TAG cells were cotransfected with 20  $\mu$ g pSX-NFAT/SEAP with 20  $\mu$ g empty vector, pEF/LAT(Wt), or pEF/LAT(Y171/191F), and the same procedure used for stimulation and SEAP assay as in (B).

FIGURE 7A shows the nucleotide sequence (SEQ ID NO:1) of human LAT.

FIGURE 7B shows the nucleotide sequence (SEQ ID NO: 2) of an alternative splice variant of human LAT.

FIGURE 7C shows the nucleotide sequence (SEQ ID NO: 3) of murine LAT.

FIGURE 7D shows the amino acid sequence (SEQ ID NO:4) of human LAT and of murine LAT (SEQ ID NO:5).

FIGURE 8A shows the nucleotide sequence (SEQ ID NO:6) of human ZAP-70 kinase.

FIGURE 8B shows the amino acid sequence (SEQ ID NO:7) of human ZAP-70 kinase.

FIGURE 9A shows the nucleotide sequence (SEQ ID NO:8) of human Syk kinase.

FIGURE 9B shows the amino acid sequence (SEQ ID NO:9) of human Syk kinase.

FIGURE 10A shows the nucleotide sequence (SEQ ID NO:10) of human Grb2.

FIGURE 10B shows the amino acid sequence (SEQ ID NO:11) of human Grb2.

FIGURE 11A shows the nucleotide sequence (SEQ ID NO:12) of human Vav.

FIGURE 11B shows the amino acid sequence (SEQ ID NO:13) of human Vav.

FIGURE 12A shows the nucleotide sequence (SEQ ID NO:14) of human Cbl.

FIGURE 12B shows the amino acid sequence (SEQ ID NO:15) of human Cbl.

FIGURE 13A shows fractions from a sucrose gradient separating a 1% Triton

lysate of unstimulated Jurkat cells. Resolved by SDS-PAGE and immunoblotted with anti-phosphotyrosine antibody (4G10).

FIGURE 13B shows fractions from a sucrose gradient separating a 1% Triton lysate of OKT3-stimulated Jurkat cells. Analyzed by SDS-PAGE and immunoblotted with anti-phosphotyrosine antibody.

FIGURE 13C shows the same membrane from (A) and (B). Was stripped and blotted with anti-PCL- $\gamma$ 1, Cbl, Vav, ZAP-70, Lck and Grb2 antibodies.

FIGURE 14 shows metabolic labeling of Wt and mutant forms of LAT with [ $^3$ H]palmitate.

FIGURE 15 shows localization of Wt and mutant LAT to the plasma membrane and glycolipid-enriched microdomains.

FIGURE 16 shows tyrosine phosphorylation of Wt and mutant LAT.

## DEFINITIONS

To facilitate understanding of the invention, a number of terms are defined below.

The abbreviations used herein are: LAT, Linker for activation of T cells. mLAT, mutant LAT (Y171/191F); PTKs, Protein tyrosine kinases; PTPS, protein tyrosine phosphatases; GST, glutathione S-transferase; TCR, T cell receptor, BCR, B cell antigen receptor; SEAP, secreted alkaline phosphatase; Tyr, tyrosine; PAGE, polyacrylamide gel electrophoresis; SDS, sodium dodecylsulfate; ITAMs, immunoreceptor tyrosine-based activation motifs; SLP-76, SH2-domain Leukocyte Protein; PLC- $\gamma$ 1, Phospholipase C-gamma1; PI3-K, phosphatidylinositol 3-kinase; IgR, Immunoglobulin receptor; SH2(C), C terminal of Src-homology 2 domain.

The terms "Lck and Fyn" refer to the protein tyrosine kinases of the src family, shown to be involved in TCR signaling.

The terms "Zap-70 and Syk" refer to the protein tyrosine kinases of the syk family, shown to be involved in TCR signaling.

The term "Vav" refers to the human oncogene derived from a locus ubiquitously expressed in hematopoietic cells.

The term "Cbl" refers to the human proto-oncogene c-cbl.

The term "Grb2" refers to the human epidermal growth factor receptor-binding protein.

The term "adaptor" or "linker protein" refers to a protein or scaffold upon which other signaling molecules assemble, for e.g. the Shc protein, Grb2, ZAP-70 etc

The term "gene" refers to a DNA sequence that comprises control and coding sequences necessary for the production of a polypeptide or precursor thereof. The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired enzymatic activity is retained.

The term "wild-type" refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designated the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications (e.g. deletions, substitutions, etc.) in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that naturally-occurring mutants can be isolated: these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

The term "transfection" as used herein refers to the introduction of a transgene into a cell. The term "transgene" as used herein refers to any nucleic acid sequence which is introduced into the genome of a cell by experimental manipulations. A transgene may be an "endogenous DNA sequence," or a "heterologous DNA sequence" (*i.e.*, "foreign DNA"). The term "endogenous DNA sequence" refers to a nucleotide sequence which is naturally found in the cell into which it is introduced so long as it does not contain some modification (*e.g.*, a point mutation, the presence of a selectable marker gene, etc.) relative to the naturally-occurring sequence. The term "heterologous DNA sequence" refers to a nucleotide sequence which is not endogenous to the cell into which it is introduced. Heterologous DNA includes a nucleotide sequence which is ligated to, or is manipulated to become ligated to, a nucleic acid sequence to which it is not ligated in nature, or to which it is ligated at a different location in nature. Heterologous DNA also includes a nucleotide sequence which is naturally found in the cell into which it is introduced and which contains some modification relative to the naturally-occurring sequence. Generally, although not necessarily, heterologous DNA encodes RNA and proteins that are not normally produced by the cell into which it is introduced. Examples of heterologous DNA include reporter genes, transcriptional and translational regulatory sequences, DNA sequences which encode selectable marker proteins (*e.g.*, proteins which confer drug resistance), *etc.* Yet another example of a heterologous

DNA includes a nucleotide sequence which encodes a ribozyme which is found in the cell into which it is introduced, and which is ligated to a promoter sequence to which it is not naturally ligated in that cell.

Transfection may be accomplished by a variety of means known to the art including calcium phosphate-DNA co-precipitation, DEAE-dextran-mediated transfection, polybrene-mediated transfection, electroporation, microinjection, liposome fusion, lipofection, protoplast fusion, retroviral infection, biolistics (*i.e.*, particle bombardment) and the like.

The term "stable transfection" or "stably transfected" refers to the introduction and integration of a transgene into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated one or more transgenes into the genomic DNA.

The term "transient transfection" or "transiently transfected" refers to the introduction of one or more transgenes into a transfected cell in the absence of integration of the transgene into the host cell's genome. The term "transient transfectant" refers to a cell which has transiently integrated one or more transgenes.

A "transgenic organism" as used herein refers to an organism in which one or more cells has been transiently transfected or stably transfected with a transgene by experimental manipulation. Transgenic organisms may be produced by several methods including the introduction of a "transgene" comprising nucleic acid (usually DNA) into an embryonic target cell or a somatic target cell of a non-human organism by way of human intervention.

The term "oligonucleotide" as used herein is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, usually more than three (3), and typically more than ten (10) and up to one hundred (100) or more (although preferably between twenty and thirty). The exact size will depend on many factors, which in turn depends on the ultimate function or use of the oligonucleotide. The oligonucleotide may be generated in any manner, including chemical synthesis, DNA replication, reverse transcription, or a combination thereof.

Because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage, an end of an oligonucleotide is referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid

sequence. even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends.

When two different, non-overlapping oligonucleotides anneal to different regions of the same linear complementary nucleic acid sequence, and the 3' end of one oligonucleotide points towards the 5' end of the other, the former may be called the "upstream" oligonucleotide and the latter the "downstream" oligonucleotide.

The term "primer" refers to an oligonucleotide which is capable of acting as a point of initiation of synthesis when placed under conditions in which primer extension is initiated. An oligonucleotide "primer" may occur naturally, as in a purified restriction digest or may be produced synthetically.

A primer is selected to be "substantially" complementary to a strand of specific sequence of the template. A primer must be sufficiently complementary to hybridize with a template strand for primer elongation to occur. A primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being substantially complementary to the strand. Non-complementary bases or longer sequences can be interspersed into the primer, provided that the primer sequence has sufficient complementarity with the sequence of the template to hybridize and thereby form a template primer complex for synthesis of the extension product of the primer.

"Hybridization" methods involve the annealing of a complementary sequence to the target nucleic acid (the sequence to be detected). The ability of two polymers of nucleic acid containing complementary sequences to find each other and anneal through base pairing interaction is a well-recognized phenomenon. The initial observations of the "hybridization" process by Marmur and Lane, *Proc. Natl. Acad. Sci. USA* 46:453 (1960) and Doty *et al.*, *Proc. Natl. Acad. Sci. USA* 46:461 (1960) have been followed by the refinement of this process into an essential tool of modern biology. Nonetheless, a number of problems have prevented the wide scale use of hybridization as a tool in human diagnostics. Among the more formidable problems are: 1) the inefficiency of hybridization; 2) the low concentration of specific target sequences in a mixture of genomic DNA; and 3) the hybridization of only partially complementary probes and targets.

With regard to efficiency, it is experimentally observed that only a fraction of the possible number of probe-target complexes are formed in a hybridization reaction. This is particularly true with short oligonucleotide probes (less than 100 bases in length). There are

three fundamental causes: a) hybridization cannot occur because of secondary and tertiary structure interactions; b) strands of DNA containing the target sequence have rehybridized (reannealed) to their complementary strand; and c) some target molecules are prevented from hybridization when they are used in hybridization formats that immobilize the target nucleic acids to a solid surface.

Even where the sequence of a probe is completely complementary to the sequence of the target, *i.e.*, the target's primary structure, the target sequence must be made accessible to the probe via rearrangements of higher-order structure. These higher-order structural rearrangements may concern either the secondary structure or tertiary structure of the molecule. Secondary structure is determined by intramolecular bonding. In the case of DNA or RNA targets this consists of hybridization within a single, continuous strand of bases (as opposed to hybridization between two different strands). Depending on the extent and position of intramolecular bonding, the probe can be displaced from the target sequence preventing hybridization.

Solution hybridization of oligonucleotide probes to denatured double-stranded DNA is further complicated by the fact that the longer complementary target strands can renature or reanneal. Again, hybridized probe is displaced by this process. This results in a low yield of hybridization (low "coverage") relative to the starting concentrations of probe and target.

With regard to low target sequence concentration, the DNA fragment containing the target sequence is usually in relatively low abundance in genomic DNA. This presents great technical difficulties; most conventional methods that use oligonucleotide probes lack the sensitivity necessary to detect hybridization at such low levels.

One attempt at a solution to the target sequence concentration problem is the amplification of the detection signal. Most often this entails placing one or more labels on an oligonucleotide probe. In the case of non-radioactive labels, even the highest affinity reagents have been found to be unsuitable for the detection of single copy genes in genomic DNA with oligonucleotide probes. See Wallace *et al.*, *Biochimie* 67:755 (1985). In the case of radioactive oligonucleotide probes, only extremely high specific activities are found to show satisfactory results. See Studencki and Wallace, *DNA* 3:1 (1984) and Studencki *et al.*, *Human Genetics* 37:42 (1985).

K.B. Mullis *et al.*, U.S. Patent Nos. 4,683,195 and 4,683,202, hereby incorporated by reference, describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. This process for



amplifying the target sequence (which can be used in conjunction with the present invention to make target molecules) consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. There can be numerous "cycles" to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to by the inventors as the "Polymerase Chain Reaction" (hereinafter PCR). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified".

The present invention specifically contemplates using primers having a portion of the nucleic acid sequence set forth in SEQ ID NO:1 in a PCR reaction to identify homologues of LAT, as well as the LAT gene in other species. Such primers are preferably less than fifty nucleotides in length (although longer primers can be used if desired).

The present invention also contemplates using probes having a portion of the nucleic acid sequence set forth in SEQ ID NO:1. The term "probe" as used herein refers to a labeled oligonucleotide which forms a duplex structure with a sequence in another nucleic acid, due to complementarity of at least one sequence in the probe with a sequence in the other nucleic acid.

In one embodiment, the present invention contemplates a polynucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO:1. As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. "Stringency" typically occurs in a range from about  $T_m - 5^\circ\text{C}$  ( $5^\circ\text{C}$  below the  $T_m$  of the probe) to about  $20^\circ\text{C}$  to  $25^\circ\text{C}$  below  $T_m$ . As will be understood by those of skill in the art, a stringent hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences. Conditions of "weak" or "low" stringency are often required with

nucleic acids that are derived from organisms that are genetically diverse, as the frequency of complementary sequences is usually low between such organisms.

The term "label" as used herein refers to any atom or molecule which can be used to provide a detectable (preferably quantifiable) signal, and which can be attached to a nucleic acid or protein. Labels may provide signals detectable by fluorescence, radioactivity, colorimetry, gravimetry, X-ray diffraction or absorption, magnetism, enzymatic activity, and the like. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes (e.g.,  $^3\text{H}$ ), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, beta-galactosidase, luciferase, alkaline phosphatase), biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance. In others, the label is part of the fusion protein, e.g. Green Fluorescent Protein (GFP).

The terms "nucleic acid substrate" and "nucleic acid template" are used herein interchangeably and refer to a nucleic acid molecule which may comprise single- or double-stranded DNA or RNA.

The term "substantially single-stranded" when used in reference to a nucleic acid substrate means that the substrate molecule exists primarily as a single strand of nucleic acid in contrast to a double-stranded substrate which exists as two strands of nucleic acid which are held together by inter-strand base pairing interactions.

The term "sequence variation" as used herein refers to differences in nucleic acid sequence between two nucleic acid templates. For example, a wild-type structural gene and a mutant form of this wild-type structural gene may vary in sequence by the presence of single base substitutions and/or deletions or insertions of one or more nucleotides. These two forms of the structural gene are said to vary in sequence from one another. A second mutant form of the structural gene may exist. This second mutant form is said to vary in sequence from both the wild-type gene and the first mutant form of the gene. It should be noted that, while the invention does not require that a comparison be made between one or more forms of a gene to detect sequence variations, such comparisons are possible using particular hybridization conditions as described in U.S. Patent No. 5,652,096, hereby incorporated by reference.

The term " $K_m$ " as used herein refers to the Michaelis-Menton constant for an enzyme and is defined as the concentration of the specific substrate at which a given enzyme yields one-half its maximum velocity in an enzyme catalyzed reaction.

The term "nucleotide analog" as used herein refers to modified or non-naturally occurring nucleotides such as 7-deaza purines (*i.e.*, 7-deaza-dATP and 7-deaza-dGTP). Nucleotide analogs include base analogs and comprise modified forms of deoxyribonucleotides as well as ribonucleotides. As used herein the term "nucleotide analog" when used in reference to substrates present in a PCR mixture refers to the use of nucleotides other than dATP, dGTP, dCTP and dTTP; thus, the use of dUTP (a naturally occurring dNTP) in a PCR would comprise the use of a nucleotide analog in the PCR. A PCR product generated using dUTP, 7-deaza-dATP, 7-deaza-dGTP or any other nucleotide analog in the reaction mixture is said to contain nucleotide analogs.

"Oligonucleotide primers matching or complementary to a gene sequence" refers to oligonucleotide primers capable of facilitating the template-dependent synthesis of single or double-stranded nucleic acids. Oligonucleotide primers matching or complementary to a gene sequence may be used in PCRs, RT-PCRs and the like.

A "consensus gene sequence" refers to a gene sequence which is derived by comparison of two or more gene sequences and which describes the nucleotides most often present in a given segment of the genes; the consensus sequence is the canonical sequence.

The term "polymorphic locus" is a locus present in a population which shows variation between members of the population (*i.e.*, the most common allele has a frequency of less than 0.95). In contrast, a "monomorphic locus" is a genetic locus at little or no variations seen between members of the population (generally taken to be a locus at which the most common allele exceeds a frequency of 0.95 in the gene pool of the population).

The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

The term "corresponds to" is used herein to mean that a polynucleotide sequence is homologous (*i.e.*, is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to

mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference sequence "TATAR" and is complementary to a reference sequence "GTATA".

The following terms are used to describe the sequence relationships between two or more polynucleotides: "reference sequence", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length LAT cDNA sequence given in a sequence listing or may comprise a complete gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions wherein a polynucleotide sequence may be compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected. The term "sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences

over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison. The reference sequence may be a subset of a larger sequence, for example, as a segment of the full-length LAT sequence.

As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine.

The term "fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion as compared to the native protein, but where the remaining amino acid sequence is identical to the corresponding positions in the amino acid sequence deduced from a full-length cDNA sequence (e.g., various fragments of LAT protein).

Fragments typically are at least 4 amino acids long, preferably at least 20 amino acids long, usually at least 50 amino acids long or longer, and span the portion of the polypeptide required for intermolecular binding of LAT.

The term "analog" as used herein refers to polypeptides which are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of the deduced amino acid sequence of LAT. The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues. Agents are evaluated for potential activity as antagonists or agonists of LAT by inclusion in screening assays described herein below.

## GENERAL DESCRIPTION OF THE INVENTION

The invention generally relates to compositions and methods for identifying and testing tyrosine kinase signaling pathway agonists and antagonists, and more particularly, methods and compositions for screening compounds and identifying compounds that will modulate the interaction of protein tyrosine kinase substrates with their intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway.

Identifying and developing new pharmaceuticals is a multibillion dollar industry in the U.S. alone. Agonists and antagonists of critical proteins in signal transduction pathways provide a promising class of targets for novel therapeutics directed to human diseases. Urgently needed are efficient methods of identifying pharmacological agents or drugs which are active at critical target points of signal transduction pathways. Methods amenable to automated, cost-effective, high throughput drug screening have immediate application in a broad range of domestic and international pharmaceutical and biotechnology drug development programs.

The present invention contemplates elements of the T cell signal transduction pathway as targets for pharmaceutical intervention in the immune system. Immunosuppression is therapeutically desirable in a wide variety of circumstances including transplantation, allergy and other forms of hypersensitivity, autoimmune diseases, infectious diseases and cancer. Cyclosporin, a widely used drug for effecting immunosuppression, is believed to act by inhibiting a calcineurin, a phosphatase which activates certain transcription factors. However, because of side effects and toxicity, clinical indications of cyclosporin (and the more recently developed FK506) are limited. Accordingly, it is desired to identify agents which specifically

target critical points in the pathway, that will be less toxic than current drugs, such as cyclosporin, which target more widely expressed molecules.

In one embodiment, the invention provides compositions and methods for identifying and testing the syk family of protein tyrosine kinase (ZAP-70 and syk) substrates. In another embodiment, the invention contemplates identifying tyrosine kinase signaling pathway agonists and antagonists, and in particular, compositions comprising LAT which is Tyr phosphorylated by ZAP-70 and/or Syk protein tyrosine kinases leading to recruitment of multiple signaling molecules.

In preferred embodiments, the invention contemplates identifying tyrosine kinase signaling pathway agonists and antagonists, and in particular, compositions comprising LAT and fragments thereof, which bind to Grb2

The description of the invention involves : A) T cell antigen receptor signaling, B) TCR-Associated Protein Tyrosine Kinases and their Substrates, C) Multimolecular Complexes in TCR signaling, D) Tyrosine Phosphatases, E) LAT as a ZAP-70 and/or Syk tyrosine kinase substrate, and F) Uses Of The Invention.

#### **A. T Cell Antigen Receptor Signaling**

The complexity of molecular associations involved in TCR signaling begins with the TCR, which is composed of 6 different polypeptide chains thought to be organized into an 8-chain structure. These polypeptides include a ligand binding heterodimer ( $\alpha\beta$  or  $\gamma\delta$ ) and the non-polymorphic CD3  $\epsilon$ ,  $\gamma$ ,  $\delta$  and TCR $\zeta$  chains. These non-polymorphic chains are required for receptor assembly, cell-surface expression and signaling. Critical regions of the cytoplasmic domains of these chains are the immunoreceptor tyrosine-based activation motifs (ITAMs), three of which are in each of the TCR $\zeta$  chains, and one in each of the CD3 chains (Cambier, 1995). Since each TCR can contain a TCR $\zeta$  dimer and two CD3 dimers ( $\epsilon\delta$ ,  $\epsilon\gamma$ ), each TCR can contain a total of ten ITAMs.

It has been well documented that these motifs are necessary and sufficient for coupling the TCR to the intracellular signaling machinery, and function by binding key signaling molecules in resting and activated T cells. Upon TCR activation the tyrosine residues within the ITAMs become phosphorylated, permitting the binding of SH2 domain-containing proteins as a consequence of the ability of SH2 domains to bind specific phosphotyrosine-containing polypeptides. Additional signaling molecules are subsequently

recruited to these newly TCR-associated proteins via SH2 or other modular interaction domains.

While a similar requirement of ITAMs for initiating signaling by the B-cell antigen receptor (BCR) and certain immunoglobulin receptors (IgR) has been shown, the TCR is notable for the sheer number of ITAMs present in a single receptor. Two different explanations for the presence of ten ITAMs within a single receptor have been proposed. The multiple ITAMs may provide the capacity to amplify the signal received by each TCR. Indeed, it has been shown that the intensity of the response to signaling by recombinant ITAM-containing polypeptide chains is dependent on the number of ITAMs present. As the T cell may be required to respond strongly in the presence of a low antigen concentration, this multiplicity of ITAMs is potentially quite important. However, recent studies offer an alternative explanation. A specificity of interaction with different ITAMs has been noted for a number of molecules including ZAP-70, PLC $\gamma$ 1, PI3-K and Shc, raising the possibility that ITAMs are bound differentially [Exley, M., Varticovsky, L., Peter, M., Sancho, J. and Terhorst, C., Association of phosphatidylinositol-3-kinase with a specific sequence of the T cell receptor chain is dependent on T cell activation, *J. Biol. Chem.* 269:15140-15146 (1994); Cambier, J. C. and Johnson, S. A., Differential binding activity of ARH1/TAM motifs, *Immunol. Lett.* 44:77-80 (1995); Isakov, N., Wange, R. L., Burgess, W. H., Watts, J. D., Aebersold, R. and Samelson, L. E., ZAP-70 binding specificity to T cell receptor tyrosine-based activation motifs: the tandem SH2 domains of ZAP-70 bind distinct tyrosine-based activation motifs with varying affinity, *J. Exp. Med.* 181:375-380 (1995); Osman, N., Turner, H., Lucas, S., Reif, K. and Cantrell, D.A., The protein interactions of the immunoglobulin receptor family tyrosine-based activation motifs present in the T cell receptor subunits and the CD3 $\epsilon$  and chains, *Eur. J. Immunol.* 26:1063-1068 (1996)]. Of course, the multiplicity of ITAMs within the TCR may actually serve both purposes.

Another important function of ITAMs may be to mediate an activation-induced association of TCR $\zeta$  and CD3 $\epsilon$  with the actin cytoskeleton. There is increasing evidence to suggest that in resting T cells a fraction of the TCR is tightly associated with the actin cytoskeleton, and that upon TCR stimulation the portion of TCR $\zeta$  and CD3 $\epsilon$  recoverable in the cytochalasin-disruptable, detergent-insoluble fraction increases [Caplan, S., Zeliger, S., Wang, L. and Baniyash, M., Cell-surface expressed T-cell antigen-receptor chain is associated with the cytoskeleton, *Proc. Natl. Acad. Sci. USA*, 92:4768-4772 (1995); Rozdzial, M. M., Malissen, B. and Finkel, T. H. (1995). Tyrosine-phosphorylated T cell receptor chain



associates with actin cytoskeleton upon activation of mature T lymphocytes. *Immunity*, 3:623-633 (1995)]. This association with the actin cytoskeleton requires the intact C-terminal ITAM of TCR $\zeta$ . Further investigation is needed to determine how these interactions impinge on TCR recycling, trafficking and signaling, and to assess the role of TCR-cytoskeletal interactions in mediating the changes in cell shape and motility patterns that can accompany T cell activation.

#### B. TCR-Associated Protein Tyrosine Kinases and their Substrates

Immediately downstream of the TCR in the signaling pathway are the TCR-associated protein tyrosine kinases (PTKs). Two families of PTKs have been shown to be involved in TCR signaling. Lck and Fyn are members of the Src family, while ZAP-70 and Syk make up another PTK family. Characteristics of these enzymes have been extensively reviewed [Bolen, J.B., Nonreceptor tyrosine protein kinases, *Oncogene*, 8:2025-2031 (1993); Peri, K. G. and Veillette, A., Tyrosine protein kinases in T lymphocytes. *Chem. Immunol.* 59:19-39 (1994)]. A primary function of the Src-family kinases is to phosphorylate key tyrosine residues within the ITAMs. An additional function of these kinases includes the phosphorylation and concurrent activation of the ZAP-70 kinase. Other substrates for these enzymes remain to be defined, though some candidates such as the receptor for IP<sub>3</sub> (providing a mechanism for Fyn to directly regulate intracellular Ca<sup>2+</sup>) have been identified [(Jayaraman *et al.*, Regulation of the Inositol 1,4,5-trisphosphate receptor by tyrosine phosphorylation, *Science* 272:1492-1494 (1996)]. It remains to be determined whether Lck and Fyn have any unique substrates, or whether their roles in signaling are redundant. Genetic experiments in which these enzymes are either deleted or over-expressed suggest unique roles for them during development, but these experiments do not address function in a mature T cell that developed in a normal environment.

The relatively recent discovery of the PTKs ZAP-70 and Syk has led to intense investigation of their function in lymphocytes. Syk is central to BCR and IgR function. Though possibly involved in T cell development and the function of certain subsets of  $\gamma\delta$  T cells, its importance to TCR signaling in mature T cells has not been demonstrated [Cheng *et al.*, Syk tyrosine kinase required for mouse viability and B-cell development, *Nature* 378:303-306 (1995); Fargnoli *et al.*, Syk mutation in Jurkat E6-derived clones results in lack of p72syk expression, *J Biol. Chem.* 270:26533-26537 (1995); Turner *et al.*, Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk, *Nature*

378:298-302 (1995)]. In contrast, study of patients with ZAP-70 deficiency and mice with a genetically engineered absence of ZAP-70 confirm the critical function of this enzyme for TCR-mediated signaling [Arpaia *et al.*, Defective T cell receptor signaling and CD8+ thymic selection in humans lacking ZAP-70 kinase, *Cell* 76:947-958 (1994); Chan *et al.*, ZAP-70 deficiency in an autosomal recessive form of severe combined immunodeficiency, *Science* 264:1599-1601 (1994); Elder *et al.*, Human severe combined immunodeficiency due to a defect in ZAP-70, a T cell tyrosine kinase, *Science* 264:1596-1599 (1994); Negishi *et al.*, Essential role for ZAP-70 in both positive and negative selection of thymocytes, *Nature* 376:435-438 (1995)].

Certain factors that regulate the activity of ZAP-70 have been identified. That activation of ZAP-70 requires a Src family kinase could be inferred from the initial description of this enzyme [Chan *et al.*, ZAP-70: a 70 kd protein-tyrosine kinase that associates with the TCR chain, *Cell* 71:649-662 (1992)]. Full enzymatic activity of ZAP-70 expressed in COS cells required co-expression of Lck or Fyn. These results, and the generally recognized ability of tyrosine phosphorylation to regulate PTK activity, lead to an investigation of the sites of tyrosine phosphorylation within activated ZAP-70. Recombinant ZAP-70, when analyzed *in vitro*, becomes tyrosine phosphorylated at a low level at Y126 and Y292 [Watts *et al.*, Identification by electrospray ionization mass spectrometry of the sites of tyrosine phosphorylation induced in activated Jurkat T cells on the protein tyrosine kinase ZAP-70, *J. Biol. Chem.* 269:29520-29529 (1994)]. However, only when assayed with added recombinant Lck does maximal tyrosine phosphorylation and maximal activity of the enzyme occur. Prominent sites of phosphorylation under these conditions are the adjacent tyrosines 492 and 493 in the kinase domain as well as tyrosines 69, 126, 178 and 292. A similar experiment using kinase-dead ZAP-70 (K369R) and the isolated kinase domain of Lck found only a single prominent site of phosphorylation in ZAP-70 at Y493 [Guanghui *et al.*, Distinct tyrosine phosphorylation sites within ZAP-70 mediate activation and negative regulation of antigen receptor function, *Mol. Cell. Biol.* (1996) In press]. ZAP-70 isolated from intact, activated T cells contains phosphate on Y492 and Y493 as well as on Y292, which lies between the second SH2 domain and the kinase domain (Watts *et al.*, Identification by electrospray ionization mass spectrometry of the sites of tyrosine phosphorylation induced in activated Jurkat T cells on the protein tyrosine kinase ZAP-70, *J. Biol. Chem.* 269:29520-29529 (1994)]. Another critical observation was that isolated proteolytic phosphopeptides purified from activated ZAP-70 contained phosphate on Y493 alone or on

both Y492 and Y493, but not on Y492 alone, suggesting that phosphorylation at Y493 precedes and is required for phosphorylation of Y492 (Chan *et al.*, Activation of ZAP-70 kinase activity by phosphorylation of tyrosine 493 is required for lymphocyte antigen receptor function. *EMBO J.* 14:2499-2508 (1995)). This has also been suggested by COS cell studies with ZAP-70 carrying Tyr to Phe mutations at either Y492 or Y493 [Wange *et al.*, Activating and inhibitory mutations in adjacent tyrosines in the kinase domain of ZAP-70. *J. Biol. Chem.* 270:18730-18733 (1995)].

These observations have led to a model describing the sequential activation of protein tyrosine kinases following TCR engagement [Weiss, A. and Littman, D. R., Signal transduction by lymphocyte antigen receptors, *Cell* 76:263-274 (1994); Cantrell, D., T Cell Antigen Receptor Signal Transduction Pathways, *Annu. Rev. Immunol.* 14: 259-274 (1996)]. Occupancy of the TCR causes the initial activation of Lck and/or Fyn leading to tyrosine phosphorylation of ITAMs. By virtue of its tandem SH2 domains, ZAP-70 then binds to these motifs [Wange *et al.*, Tandem SH2 domains of ZAP-70 bind to T cell antigen receptor and CD3 from activated Jurkat T cells, *J. Biol. Chem.* 268:19797-19801 (1993); Iwashima *et al.*, Sequential interactions of the TCR with two distinct cytoplasmic tyrosine kinases, *Science* 263:1136-1139 (1994)]. Recruitment of ZAP-70 to the ITAMs is required for activation of its kinase activity and for T cell activation, since agents that block recruitment prevent these events [Wange *et al.*, F2(Pmp)2-TAM 3, a novel competitive inhibitor of the binding of ZAP-70 to the T cell antigen receptor, blocks early T cell signaling, *J. Biol. Chem.* 270:944-948 (1995); Qian *et al.*, Dominant-negative Zeta-associated Protein 70 Inhibits T Cell Antigen Receptor Signaling, *J. Exp. Med.* 183:611-620 (1996)]. This sequence of events can be detected in cultured cell lines and in peripheral blood lymphocytes. However, in thymocytes and lymph node T cells, one can detect inactive, non-phosphorylated ZAP-70 bound to the basally tyrosine-phosphorylated TCR in the absence of activation [van Oers *et al.*, Constitutive tyrosine phosphorylation of the T-cell receptor (TCR) subunit: regulation of TCR-associated protein tyrosine kinase activity by TCR, *Mol. Cell. Biol.* 13:5771-5780 (1993); Wiest *et al.*, TCR activation of ZAP-70 is impaired in CD4+CD8+ thymocytes as a consequence of intrathymic interactions that diminish available p56lck. *Immunity* 4:495-504 (1996)]. The phosphorylation of ZAP-70 on Y493 by a Src-family kinase results in the activation of ZAP-70 kinase activity [Chan *et al.*, Activation of ZAP-70 kinase activity by phosphorylation of tyrosine 493 is required for lymphocyte antigen receptor function. *EMBO J.* 14:2499-2508 (1995); Wange *et al.*, Activating and inhibitory mutations in adjacent

tyrosines in the kinase domain of ZAP-70, *J. Biol. Chem.* 270:18730-18733 (1995)].

Although this site shares sequence similarity with the so-called autophosphorylation sites defined on other PTKs, for ZAP-70 it is more accurate to refer to this as a transphosphorylation/activation site, since ZAP-70 is unable to phosphorylate this site, and phosphorylation of this site is required for full kinase activity. The mechanism of subsequent tyrosine phosphorylation events within the kinase domain at Y492, outside this region at Y292, and perhaps at other sites is not defined, but is likely to be due to autophosphorylation or transphosphorylation by other activated ZAP-70 molecules on adjacent ITAMs, although an additional role for a heterologous kinase has not been ruled out.

Clearly the role of phosphorylation of Y493 is to activate the kinase activity of ZAP-70; however, the roles played by phosphorylation of tyrosines 492 and 292 are less clear. In contrast to what is seen in Y493F mutants, the mutation of Y492 to Phe results in increased activity in ZAP-70 isolated from transfected COS cells and enhanced BCR signaling in Syk<sup>-/-</sup> B cells reconstituted with this ZAP-70 mutant [Wange *et al.*, Activating and inhibitory mutations in adjacent tyrosines in the kinase domain of ZAP-70, *J. Biol. Chem.* 270:18730-18733 (1995); Guanghai *et al.*, Distinct tyrosine phosphorylation sites within ZAP-70 mediate activation and negative regulation of antigen receptor function, *Mol. Cell. Biol.*, In press (1996)]. Tyrosine Y492 may therefore serve a negative regulatory function in ZAP-70 when phosphorylated. Alternatively, mutation of Y492 to Phe may cause a loss of hydrogen bonding capacity that maintains the kinase in an inactive conformation when unphosphorylated [Wange *et al.*, Activating and inhibitory mutations in adjacent tyrosines in the kinase domain of ZAP-70, *J. Biol. Chem.* 270:18730-18733 (1995)]. Mutation of Y292 to Phe, unlike the Y492F mutation, has no effect on the activity of purified ZAP-70, but, like the Y492F mutation, results in a hyperactivated state upon BCR stimulation in the Syk<sup>-/-</sup> chicken B cells [Guanghai *et al.*, Distinct tyrosine phosphorylation sites within ZAP-70 mediate activation and negative regulation of antigen receptor function, *Mol. Cell. Biol.*, In press (1996)]. This is consistent with phosphorylation of Y292 serving a role as a binding site for regulatory molecules, such as SH2 domain-bearing protein tyrosine phosphatases (see below).

Recent reports have identified yet another class of PTKs, the Itk/Btk/Tec family, that may be involved in TCR signaling [reviewed in Desiderio, S. and Siliciano, J. D., The Itk/Btk/Tec Family of Protein-Tyrosine Kinases, *Chem. Immunol.* 59:191-208 (1994)]. This family of PTKs is characterized by having an N-terminal Plextrin Homology (PH) domain.

and an SH2 and SH3 domain, in addition to the C-terminal kinase domain. This family of PTKs was first shown to be important in hematopoietic cell function when it was discovered that mutations in the Btk gene at the XLA locus cause X-linked agammaglobulinemia in man. Thus Btk, which is expressed predominantly in B lymphoid and myelomonocytic lineages, is required for normal B cell function. Itk is predominantly expressed in T cells. Itk<sup>-/-</sup> mice produce fewer thymocytes, and mature T cells isolated from these mice proliferate poorly in response to TCR stimulation, but respond normally to phorbol ester plus ionomycin [Liao, X. C. and Littman, D., Altered T cell receptor signaling and disrupted T cell development in mice lacking Itk. *Immunity*, 3:757-769 (1995)]. In Jurkat T cells, TCR cross-linking rapidly and transiently tyrosine-phosphorylates Itk, resulting in increased kinase activity in anti-Itk immunoprecipitates (Gibson *et al.*, The EMT/ITK/TSK (EMT) tyrosine kinase is activated during TCR signaling. *J. Immunol.* 156:2716-2722 (1996)]. The precise role of Itk in TCR signaling remains to be determined.

Characterization of pathways downstream of the PTKs depends on identification of their substrates. Detection of protein tyrosine phosphorylation by immunoblotting with specific anti-phosphotyrosine antibodies has revealed a number of kinase targets [Peri, K. G. and Veillette, A., Tyrosine protein kinases in T lymphocytes. *Chem. Immunol.* 59:19-39 (1994)]. In this fashion PLC $\gamma$ 1 was the first substrate, other than the TCR subunits, to be identified in T cells. Tyrosine phosphorylation of this enzyme is critical to its activation. Vav, a hematopoietic cell-specific proto-oncogene, and more recently, the proto-oncogene Cbl were also identified as being PTK substrates in T cells [Bustelo *et al.*, Product of the Vav protooncogene defines a new class of tyrosine protein kinase substrates. *Nature* 356: 68-74 (1992); Donovan *et al.*, The protein product of the c-cbl protooncogene is the 120-kDa tyrosine-phosphorylated protein in Jurkat cells activated via the T cell antigen receptor. *J. Biol. Chem.* 269:22921-22924 (1994)]. SH2-domain Leukocyte Protein (SLP-76), a prominent substrate, was recently isolated by virtue of an interaction with the Grb2 linker protein as described below [Jackman *et al.*, Molecular cloning of SLP-76, a 76-kDa tyrosine phosphoprotein associated with Grb2 in T cells, *J. Biol. Chem.* 270:7029-7032 (1995)]. More difficult than identifying the proteins that become tyrosine phosphorylated upon TCR stimulation is determining which PTKs are directly responsible for a given phosphorylation event. As stated above it is clear that ITAM phosphorylation is a function of the Src-family kinases, as is phosphorylation of Y493 of ZAP-70. ZAP-70, at least *in vitro*, does not phosphorylate these sites.

Substrates of ZAP-70 have been more difficult to identify. Based on previous study of Syk, it has been shown that ZAP-70 can phosphorylate the cytoplasmic domain of erythrocyte band 3 (cdb3) and tubulin *in vitro* (Wange et al., 1995b; Isakov et al., 1996). Evidence that cdb3 and tubulin are genuine *in vivo* substrates of Syk has also been presented [Harrison *et al.*, Phosphorylation of human erythrocyte band 3 by endogenous p72syk, *J. Biol. Chem.* 269:955-959 (1994); Peters *et al.*, Syk, activated by cross-linking the B-cell antigen receptor, localizes to the cytosol where it interacts with and phosphorylates alpha-tubulin on tyrosine, *J. Biol. Chem.* 271:4755-4762 (1996)]. Whether tubulin is an *in vivo* substrate of ZAP-70 remains to be determined; however, it is interesting that tubulin is tyrosine phosphorylated in activated T cells, and both ZAP-70 and Vav have been found in association with tubulin [Ley *et al.*, Tyrosine phosphorylation of tubulin in human T lymphocytes, *Eur. J. Immunol.*, 24:99-106 (1994); Huby *et al.*, Interactions between the Protein-tyrosine Kinase ZAP-70, the Proto-oncoprotein Vav, and Tubulin in Jurkat T Cells, *J. Biol. Chem.*, 270:30241-30244 (1995)]. These findings further support the contention that cytoskeletal components may play an integral part in the early signal transduction steps. Of the molecules involved in TCR signal transduction that undergo tyrosine phosphorylation upon TCR engagement, only SLP-76 has been shown to be a substrate of ZAP-70 [Wardenburg *et al.*, Phosphorylation of SLP-76 by the ZAP-70 protein tyrosine kinase is required for T cell receptor function, *J. Biol. Chem.*, in press (1996)]. Tyrosine phosphorylation of SLP-76 mediates association with Vav (see below) (Wu *et al.*, Vav and SLP-76 interact and functionally cooperate in IL-2 gene activation, *Immunity* 4:593-602 (1996)]. Of note is that sites likely to be phosphorylated in these three proteins, all share the sequence Asp-Tyr-Glu. Interestingly, the hematopoietic cell specific protein HS1, which has two Asp-Tyr-Glu sites, has been shown to be a substrate for ZAP-70 when HS1, ZAP-70 and Fyn are co-expressed in COS cells (Fusaki *et al.*, Physical and functional interactions of protein tyrosine kinases, p59fyn and ZAP-70, in T cell signaling, *J. Immunol.* 156:1369-1377 (1996)]. The function of HS1 remains unknown.

### C. Multimolecular Complexes in TCR signaling

While it is not intended that the present invention be limited to any particular mechanism of receptor-mediated signal transduction, it is believed that receptor activation leads to assembly of multimolecular complexes on the cytoplasmic domain of platelet-derived and epidermal growth factor receptors. These complexes comprise of multiple signaling molecules that often bind to the receptor by interaction at sites of tyrosine phosphorylation.

Many of these signaling molecules are themselves modular, containing variable numbers of interaction domains such as (1) SH2, (2) PTB, (3) SH3 and (4) PH which bind, respectively to (1) pY (in the context of specific residues C-terminal of pY), (2) pY (in the context of specific residues N-terminal of pY), (3) proline-rich regions and (4) phosphatidylinositol (4,5) biphosphate, inositol phosphates and certain proteins such as G protein  $\beta\gamma$  subunits. PH domains are thus thought to target some proteins to the plasma membrane. It is believed that, this modular architecture permits the assembly of large multimolecular complexes, as each signaling protein may bind to several different signaling proteins, which may themselves associate with a whole host of additional signaling molecules.

Proteins with the potential to become involved in multimolecular signaling complexes at the TCR include the receptor-proximal kinases themselves. The Src-family kinases contain an SH2, an SH3 and a unique N-terminal domain, all of which can be involved in protein-protein interactions [Bolen, J.B. . Nonreceptor tyrosine protein kinases. *Oncogene* 8: 2025-2031 (1993); Peri, K. G. and Veillette, A., Tyrosine protein kinases in T lymphocytes. *C'hem.Immunol.* 59:19-39 (1994)]. In addition, lipid modification of consensus N-terminal sites stabilizes association of these PTKs with cellular membranes. The unique N-terminal domain of Lck binds to the co-receptor molecules CD4 or CD8, while the comparable region in Fyn binds to non-phosphorylated ITAMs. The SH2 and SH3 domains of Src-family PTKs form an intramolecular association with the tyrosine-phosphorylated C-terminal tail, inhibiting their kinase activity. The phosphorylation status of this negative regulatory tyrosine is controlled by the competing activities of the Csk PTK and the CD45 protein tyrosine phosphatase (PTP). In addition to activating the kinase activity, dephosphorylation of this residue by CD45 also increases the availability of the SH2 and SH3 domains for interaction with other proteins (eg. Cbl, phosphatidylinositol 3-kinase [PI3-K] and ZAP-70) [Donovan *et al.*, The protein product of the c-cbl protooncogene is the 120-kDa tyrosine-phosphorylated protein in Jurkat cells activated via the T cell antigen receptor, *J. Biol. Chem.* 269:22921-22924 (1994); Vogel, L. B. and Fujita, D. J., The SH3 domain of p56lck is involved in binding to phosphatidylinositol 3'-kinase from T lymphocytes, *Mol. Cell. Biol.*, 13:7408-7417 (1993); Thome *et al.*, Syk and ZAP-70 mediate recruitment of p56lck/CD4 to the activated T cell receptor/CD3/ complex, *J. Exp. Med.*, 181:1997-2006 (1995)].

ZAP-70 is also capable of forming multimolecular complexes through its two SH2 domains and via tyrosine-phosphorylated SH2 domain acceptor sites. The only known function of the SH2 domains of ZAP-70 is to target the kinase to the two phosphorylated

tyrosines of the ITAM [Wange *et al.*, Tandem SH2 domains of ZAP-70 bind to T cell antigen receptor and CD3 from activated Jurkat T cells, *J. Biol. Chem.* 268:19797-19801 (1993); Iwashima *et al.*, Sequential interactions of the TCR with two distinct cytoplasmic tyrosine kinases. *Science* 263:1136-1139 (1994)]. It is possible however that these SH2 domains have other targets when not engaged to the activated TCR, but this seems unlikely given the unique structure of the tandem SH2 domain of ZAP-70, which only permits high affinity binding to polypeptides possessing two phosphotyrosines within a prescribed distance, as is found in ITAMs [Hatada *et al.*, Molecular basis for interaction of the protein tyrosine kinase ZAP-70 with the T-cell receptor, *Nature* 377:32-38 (1995)]. It is important to note that, unlike Syk, ZAP-70 is not activated by the binding of its tandem SH2 domains to ITAMs [Shiue *et al.*, Syk is activated by phosphotyrosine-containing peptides representing the tyrosine-based activation motifs of the high affinity receptor for IgE, *J. Biol. Chem.* 270:10498-10502 (1995); Neumeister *et al.*, Binding of ZAP-70 to phosphorylated T-cell receptor and enhances its autophosphorylation and generates specific binding sites for SH2 domain-containing proteins. *Mol. Cell. Biol.* 15:3171-3178 (1995); Isakov *et al.*, Purification and characterization of human ZAP-70 protein tyrosine kinase from a baculovirus expression system, *J. Biol. Chem.* 271:15753-15761 (1996)].

In addition to its role as an enzyme responsible for protein tyrosine phosphorylation, ZAP-70 appears also to function as an adaptor protein or scaffold upon which other signaling molecules assemble [Neumeister *et al.*, Binding of ZAP-70 to phosphorylated T-cell receptor and enhances its autophosphorylation and generates specific binding sites for SH2 domain-containing proteins, *Mol. Cell. Biol.* 15:3171-3178 (1995)]. Interestingly, Lck, Vav and Cbl have all been found in association with ZAP-70 [Thome *et al.*, Syk and ZAP-70 mediate recruitment of p56lck/CD4 to the activated T cell receptor/CD3/ complex. *J. Exp. Med.* 181:1997-2006 (1995); Katzav *et al.*, The protein tyrosine kinase ZAP-70 can associate with the SH2 domain of proto-vav, *J. Biol. Chem.* 269:32579-32585 (1994); Fournel *et al.*, Association of tyrosine protein kinase ZAP-70 with the protooncogene product p120c-cbl in T lymphocytes. *J. Exp. Med.* 183:301-306 (1996)], and, at least for Lck and Vav, this association appears to involve the SH2 domains of these two proteins binding to phosphorylated tyrosines on ZAP-70. As mentioned previously tyrosines 69, 126, 178, 292, 492 and 493 have all been shown to be capable of accepting phosphate in an *in vitro* kinase reaction with Lck [Watts *et al.*, Identification by electrospray ionization mass spectrometry of the sites of tyrosine phosphorylation induced in activated Jurkat T cells on the protein



tyrosine kinase ZAP-70. *J. Biol. Chem.* 269:29520-29529 (1994)]. It has been suggested that two other tyrosines (315 and 319) may be phosphorylatable, and phosphorylated peptides cognate for this region can disrupt the association of Vav with ZAP-70 (Katzav *et al.*. The protein tyrosine kinase ZAP-70 can associate with the SH2 domain of proto-vav. *J. Biol. Chem.* 269: 32579-32585 (994)] While the functional consequences of the ZAP-70 association with these signaling molecules remains to be determined, one can speculate that these associations ensure their efficient phosphorylation by ZAP-70 or associated PTKs, such as Lck. In addition, the association of Lck with tyrosine-phosphorylated ZAP-70 has been suggested to be required for recruitment of Lck-CD4 to the TCR [Thome *et al.*, Syk and ZAP-70 mediate recruitment of p56lck/CD4 to the activated T cell receptor/CD3/ complex, *J. Exp. Med.* 181:1997-2006 (1995)], thereby enhancing the assembly of a fully effective signaling complex.

The adaptor protein Grb2 is an example of a modular, so-called, linker protein, consisting entirely of a central SH2 domain flanked by two SH3 domains. In a number of growth factor receptor tyrosine kinase systems, Grb2 has been shown to couple these receptors to the Ras activator protein SOS [Downward, J., The GRB2/Sem-5 adaptor protein, *FEBS Lett.* 338: 113-117 (1994)] As the Ras pathway was known to be required for lymphokine gene activation, a number of investigators sought to determine whether Grb2 was linked to this pathway in T cells. The Grb2-SOS interaction was indeed documented and SOS has been shown to be involved in Ras activation in T cells [Downward, J., The GRB2/Sem-5 adaptor protein, *FEBS Lett.* 338: 113-117 (1994); Holsinger *et al.*, Signal transduction in T lymphocytes using a conditional allele of Sos, *Proc. Natl. Acad. Sci. USA* 92:9810-9814 (1995)]. While, Shc, another linker protein, has been shown to link Grb2 to certain growth factor receptors, its role in TCR signaling remains controversial [Ravichandran *et al.*, Interaction of Shc with Grb2 regulates association of Grb2 with mSOS, *Mol. Cell. Biol.* 15:593-600 (1995); Osman *et al.*, A comparison of the interaction of Shc and the tyrosine kinase ZAP-70 with the T cell antigen receptor chain tyrosine-based activation motif, *J. Biol. Chem.* 270:13981-13986 (1995)].

In addition to the anticipated role of Grb2 in Ras activation, Grb2 was also found to bind several of the most prominent substrates of the TCR-associated PTKs [Motto *et al.*, In vivo association of Grb2 with pp116, a substrate of the T cell antigen receptor-activated protein tyrosine kinase. *J. Biol. Chem.* 269:21608-21613 (1994); Buday *et al.*, A complex of Grb2 adaptor protein, Sos exchange factor, and a 36-kDa membrane-bound tyrosine

phosphoprotein is implicated in ras activation in T cells. *J. Biol. Chem.* 269:9019-9023 (1994); Reif *et al.*, SH3 domains of the adapter molecule Grb2 compete with two proteins in T cells: the guanine nucleotide exchange protein SOS and a 75-kDa protein that is a substrate for T cell antigen receptor-activated tyrosine kinases. *J. Biol. Chem.* 269:14081-14087 (1994)]. A 120kD protein that binds to Grb2 has been shown to be the proto-oncogene Cbl [Donovan *et al.*, The protein product of the c-cbl protooncogene is the 120-kDa tyrosine-phosphorylated protein in Jurkat cells activated via the T cell antigen receptor. *J. Biol. Chem.* 269:22921-22924 (1994); Fukazawa *et al.*, T cell activation-dependent association between the p85 subunit of the phosphatidylinositol 3-kinase and Grb2/phospholipase C- $\gamma$ 1-binding phosphotyrosyl protein. *J. Biol. Chem.* 270:36-38, 20177-20182 (1995); Meisner *et al.*, Interaction of Cbl with Grb2 and Phosphatidylinositol-3'-Kinase in Activated Jurkat Cells. *Mol. Cell. Biol.* 15:3571-3578 (1995)]. The Grb2-Cbl association is mediated via the N-terminal SH3 domain of Grb2, and is observed in resting and activated T cells. Affinity purification with a Grb2 fusion protein was used to isolate a 76kD protein, SLP-76, which binds to the C-terminal SH3 domain of Grb2 [Jackman *et al.*, Molecular cloning of SLP-76, a 76-kDa tyrosine phosphoprotein associated with Grb2 in T cells, *J. Biol. Chem.* 270, 7029-7032 (1995). The identity of pp36, a 36kD substrate, that binds to the SH2 domain of Grb2, has been identified to be "LAT", the composition claimed in the present invention. This protein plays a critical role in TCR signaling, as in addition to binding Grb2, it has also been found in association with PLC $\gamma$ 1 and PI3-K [Cantrell, D., T Cell Antigen Receptor Signal Transduction Pathways, *Annu. Rev. Immunol.* 14: 259-274 (1996) and also see experimental section]. The tight membrane localization of pp36 provides a possible mechanism for recruitment of Grb2, PLC $\gamma$ 1 and PI3-K to the plasma membrane. Grb2-SOS activates the Ras-pathway when localized to the plasma membrane [Downward, J., The GRB2/Sem-5 adaptor protein, *FEBS Lett.* 338: 113-117 (1994); Cantrell, D., T Cell Antigen Receptor Signal Transduction Pathways, *Annu. Rev. Immunol.* 14: 259-274 (1996)], while membrane localization of PLC $\gamma$ 1 and PI3-K permits phosphorylation of these proteins by membrane-associated PTKs, and provides access to lipid substrates.

The binding of SOS and Cbl to Grb2 is mutually exclusive, as proline-rich regions in these proteins compete for the same binding site [Meisner *et al.*, Interaction of Cbl with Grb2 and Phosphatidylinositol-3'-Kinase in Activated Jurkat Cells, *Mol. Cell. Biol.* 15:3571-3578 (1995)]. An interesting area of investigation is whether or not this competition plays a role in TCR signaling. It is also possible that Grb2 could mediate a complex between either pp36

and/or SLP-76 with Cbl or SOS. An additional consequence of formation of the Grb2-Cbl complex could depend on the ability of Cbl, itself, to serve as an adaptor protein. Recent studies demonstrate that Cbl can be found in complex with PI3-K, ZAP-70, 14.3.3 $\tau$  and CrkL [Meisner *et al.*, Interaction of Cbl with Grb2 and Phosphatidylinositol-3'-Kinase in Activated Jurkat Cells, *Mol. Cell. Biol.* 15:3571-3578 (1995); Fukazawa *et al.*, T cell activation-dependent association between the p85 subunit of the phosphatidyl 3-kinase and Grb2/phospholipase C- $\gamma$ 1-binding phosphotyrosyl protein, *J. Biol. Chem.* 270:36-38, 20177-20182 (1995); Fournel *et al.*, Association of tyrosine protein kinase ZAP-70 with the protooncogene product p120c-cbl in T lymphocytes, *J. Exp. Med.* 183:301-306 (1996); Liu *et al.*, Activation-modulated association of 14-3-3 proteins with Cbl in T cells, *J. Biol. Chem.* 271, 14591-14595 (1996); Reedquist *et al.*, Stimulation through the T cell receptor induces Cbl association with Crk proteins and guanine nucleotide exchange protein C3G, *J. Biol. Chem.* 271:8435-8442 (1996)]. CrkL is another adapter protein that in turn interacts with C3G, a nucleotide exchange factor for members of the Rac/Rho family of small G proteins.

Study of Vav and SLP-76 has led to the most significant recent insights into TCR signal transduction mechanisms. Vav, first identified as a transforming oncogene when truncated, consists of a number of interaction modules, 2 SH3 domains, one SH2 domain and one PH domain, along with regions sharing homology with Rac/Rho guanine nucleotide exchange proteins [Bustelo *et al.*, Product of the Vav protooncogene defines a new class of tyrosine protein kinase substrates, *Nature* 356:68-74 (1992)]. Over-expression of this protein in Jurkat T cells results in enhanced basal activation of IL-2 promoters and further enhances the response to TCR signaling (Wu *et al.*, Vav and SLP-76 interact and functionally cooperate in IL-2 gene activation, *Immunity* 4:593-602 (1995)]. Its proximal position in the TCR pathway was established by the observation that dominant negative Ras or Raf could block the Vav-mediated enhancement of TCR signaling. The position of Vav in the signaling cascade has been further delineated by the observation that the activity of Vav requires active PTKs and is not functional in a T cell line lacking Lck. Similar studies with SLP-76 over-expression also demonstrated an enhanced response to TCR engagement although without an increase in basal levels of activation (Motto *et al.*, Implications of the GRB2-associated phosphoprotein SLP-76 in T cell receptor-mediated interleukin 2 production, *J. Exp. Med.* 183:1937-1943 (1996)]. Over-expression of both Vav and SLP-76 causes a synergistic induction of basal and TCR-stimulated NFAT and IL-2 promoter activation [Wu *et*

al., Vav and SLP-76 interact and functionally cooperate in IL-2 gene activation. *Immunity* 4:593-602 (1996)].

How these two signaling molecules ultimately affect transcriptional activity is unknown, but it has been shown that a specific interaction between Vav and SLP-76 is required for this activity [Wu *et al.*, Vav and SLP-76 interact and functionally cooperate in IL-2 gene activation. *Immunity* 4:593-602 (1996)]. This association requires the SH2 domain of Vav and tyrosine phosphorylation of SLP-76. This association is required for the subsequent tyrosine phosphorylation of Vav, as crippling the SH2 domain of Vav prevents its tyrosine phosphorylation. Similarly, enhanced signaling does not occur if critical tyrosine residues (112, 128, 145) in SLP-76, presumably the binding sites for the Vav SH2 domain, are mutated to Phe. SLP-76, in turn binds two tyrosine-phosphorylated proteins of 130 (SLAP) and 62kD. An inactivating mutation within the SLP-76 SH2 domain prevents association of these proteins and significantly blocks the stimulatory effect of SLP-76 over-expression. Clearly Vav and SLP-76 have critical functional effects and the complex of these two proteins with others (eg. pp130 and pp62) is central to TCR function.

#### D. Tyrosine Phosphatases

The addition of phosphate to tyrosine residues catalyzed by tyrosine kinases induces many binding interactions as described. Regulation of these events by tyrosine phosphatases has equal significance both in initiating and quenching TCR signaling pathways [for review see McFarland *et al.*, Protein tyrosine phosphatases involved in lymphocyte signal transduction, *Chem. Immunol.* 59:40-61 (1994)]. Activation of Src family kinases by CD45 has already been mentioned. Recent evidence for interaction of the SHP-1 PTP with ZAP-70 suggests that regulation of this kinase or its substrates by SHP-1 may also be significant [Plas *et al.*, Direct regulation of ZAP-70 by SHP-1 in T cell antigen receptor signaling. *Science* 272:1173-1176 (1996). This association is mediated by the SH2 domains of SHP-1, which bind to tyrosine phosphorylated ZAP-70 after TCR engagement. Engagement of the SH2 domains of SHP-1 by ZAP-70 stimulates the phosphatase activity with a consequent decrease in net ZAP-70 kinase activity. The functional significance of this interaction can be demonstrated by overexpression of wildtype SHP-1, which decreases IL-2 production in response to TCR stimulation, or conversely by overexpression of catalytically inactive SHP-1 (C453S), which has the opposite effect of enhancing IL-2 production. Other, as yet, unidentified PTPs are likely to have functional significance, and substrates of known PTPs

remain to be defined. It is also interesting that SHP-1 has been found to associate with Vav in activated splenocytes and the EL4 T cell lymphoma [Kon-Kozlowski, M. et al., The tyrosine phosphatase PTP1C associates with Vav, Grb2, and mSos1 in hematopoietic cells. *J. Biol. Chem.* 271:3856-3862 (1996)]. This association appears to involve the SH2 and flanking SH3 domains of Vav. As Vav and ZAP-70, and ZAP-70 and SHP-1 have also been shown to associate, determining the actual nature of the associations between these proteins will be required to more fully understand these signaling events. Another PTP that probably plays a role in down-regulating the TCR signal is SHP-2, which has been found to associate with the activation-upregulated T cell surface protein CTLA-4. Fyn, Lck and ZAP-70 are all hyperphosphorylated in CTLA-4<sup>-/-</sup> mice [Marengere et al., Regulation of T cell receptor signaling by tyrosine phosphatase SYP association with CTLA-4. *Science* 272:1170-1173 (1996)].


#### E. LAT as a ZAP-70 and/or Syk tyrosine kinase substrate

In the present invention, LAT, one of the most prominently tyrosine phosphorylated protein following TCR engagement has been identified. Deduced amino acid sequence identifies a novel integral membrane protein containing multiple potential tyrosine phosphorylation sites. This 36-38kD protein is capable of binding SH2 domains of Grb2, phospholipase C- $\gamma$ 1 and the p85 subunit of phosphoinositide 3-kinase and may play a central role as a molecule downstream of PTKs capable of binding critical PTK substrates and effector molecules. In the present invention, this protein is shown to be phosphorylated by ZAP-70 and/or Syk protein tyrosine kinases leading to recruitment of multiple signaling molecules. Its function is demonstrated by inhibition of T cell activation following overexpression of a mutant form lacking critical tyrosine residues (For details see experimental section ), providing evidence that LAT plays an important role in linking the TCR to cellular activation. Therefore, the molecule has been named: LAT for linker for activation of T cells (See Schematic A).

#### F. Uses Of The Invention

The invention provides compositions and methods for identifying and testing tyrosine kinase ZAP-70 and/or Syk substrates and ZAP-70 and/or Syk kinase signaling pathway agonists and antagonists, and in particular, compositions comprising LAT (and fragments

LAT is a central molecule that links the TCR to cellular activation



Schematic A

thereof) which upon activation/tyrosine phosphorylation binds to ZAP-70 and/or Syk kinase and also to Grb2, PLC- $\gamma$ -1, the p85 subunit of PI-3 kinase, and Cbl, Vav, SLP-76 and/or other critical signaling molecules, either directly or indirectly upon T cell activation.

The polypeptide LAT and fragments thereof, may have one or more LAT-specific binding affinities for particular ligands such as Grb2, PLC- $\gamma$ -1, the p85 subunit of PI-3 kinase, and other critical signaling molecules, including the ability to specifically bind at least one natural human intracellular LAT-specific binding target or a LAT-specific binding agent such as a LAT-specific antibody or a LAT-specific binding agent identified in assays such as described below. Accordingly, the specificity of a LAT fragment for specific binding agents is confirmed by ensuring non-crossreactivity with other ZAP-70 and/or Syk substrates. Furthermore, preferred LAT fragments are capable of eliciting an antibody capable of distinguishing LAT from other LAT homologues.

### 1. Antibody Generation

Both polyclonal and monoclonal antibodies are obtainable by immunization with LAT, LAT fragments, mutant LAT, mutant LAT fragments, and LAT/LAT binding ligand complexes and either type is utilizable for immunoassays (as well as therapy). Polyclonal sera are readily prepared by injection of a suitable laboratory animal with an effective amount of the purified peptide, collecting serum from the animal, and isolating specific sera by any of the known immunoabsorbent techniques. Antibodies produced by this method are utilizable in virtually any type of immunoassay (see below).

The use of monoclonal antibodies directed to LAT, LAT fragments, mutant LAT, mutant LAT fragments, and LAT/LAT binding ligand complexes is particularly preferred because of the ability to produce them in large quantities and the homogeneity of the product. The preparation of hybridoma cell lines for monoclonal antibody production derived by fusing an immortal cell line and lymphocytes sensitized against the immunogenic preparation can be done by techniques which are well known to those who are skilled in the art. (See, for example Douillard and Hoffman, Basic Facts about Hybridomas, in Compendium of Immunology Vol II, ed. by Schwartz, 1981; Kohler and Milstein, *Nature* 256: 495-499, 1975; *European Journal of Immunology* 6: 511-519, 1976).

Unlike preparation of polyclonal sera, the choice of animal is dependent on the availability of appropriate immortal lines capable of fusing with lymphocytes. Mouse and rat have been the animals of choice in hybridoma technology and are preferably used. Humans

can also be utilized as sources for sensitized lymphocytes if appropriate immortalized human (or nonhuman) cell lines are available. For the purpose of the present invention, the animal of choice may be injected with an antigenic amount, for example, from about 0.1 mg to about 20 mg of the peptide or antigenic parts thereof. Usually the injecting material is emulsified in Freund's complete adjuvant. Boosting injections may also be required. The detection of antibody production can be carried out by testing the antisera with appropriately labelled antigen. Lymphocytes can be obtained by removing the spleen of lymph nodes of sensitized animals in a sterile fashion and carrying out fusion. Alternatively, lymphocytes can be stimulated or immunized *in vitro*, as described, for example, in Reading, *Journal of Immunological Methods* 53: 261-291, 1982.

A number of cell lines suitable for fusion have been developed and the choice of any particular line for hybridization protocols is directed by any one of a number of criteria such as speed, uniformity of growth characteristics, deficiency of its metabolism for a component of the growth medium, and potential for good fusion frequency.

Intraspecies hybrids, particularly between like strains, work better than interspecies fusions. Several cell lines are available, including mutants selected for the loss of ability to secrete myeloma immunoglobulin.

Cell fusion can be induced either by virus, such as Epstein-Barr or Sendai virus, or polyethylene glycol. Polyethylene glycol (PEG) is the most efficacious agent for the fusion of mammalian somatic cells. PEG itself may be toxic for cells and various concentrations should be tested for effects on viability before attempting fusion. The molecular weight range of PEG may be varied from 1000 to 6000. It gives best results when diluted to from about 20% to about 70% (w/w) in saline or serum-free medium. Exposure to PEG at 37°C for about 30 seconds is preferred in the present case, utilizing murine cells. Extremes of temperature (i.e., about 45°C) are avoided, and preincubation of each component of the fusion system at 37°C prior to fusion can be useful. The ratio between lymphocytes and malignant cells is optimized to avoid cell fusion among spleen cells and a range of from about 1:1 to about 1:10 is commonly used.

The successfully fused cells can be separated from the myeloma line by any technique known by the art. The most common and preferred method is to choose a malignant line which is Hypoxanthine Guanine Phosphoribosyl Transferase (HGPRT) deficient, which will not grow in an aminopterin-containing medium used to allow only growth of hybrids and which is generally composed of hypoxanthine  $1 \times 10^{-4} \text{M}$ , aminopterin  $1 \times 10^{-5} \text{M}$ , and thymidine



3x10<sup>-5</sup>M. commonly known as the HAT medium. The fusion mixture can be grown in the HAT-containing culture medium immediately after the fusion 24 hours later. The feeding schedules usually entail maintenance in HAT medium for two weeks and then feeding with either regular culture medium or hypoxanthine, thymidine-containing medium.

5 The growing colonies are then tested for the presence of antibodies that recognize the antigenic preparation. Detection of hybridoma antibodies can be performed using an assay where the antigen is bound to a solid support and allowed to react to hybridoma supernatants containing putative antibodies. The presence of antibodies may be detected by "sandwich" techniques using a variety of indicators. Most of the common methods are sufficiently  
10 sensitive for use in the range of antibody concentrations secreted during hybrid growth.

Cloning of hybrids can be carried out after 21-23 days of cell growth in selected medium. Cloning can be performed by cell limiting dilution in fluid phase or by directly selecting single cells growing in semi-solid agarose. For limiting dilution, cell suspensions are diluted serially to yield a statistical probability of having only one cell per well. For the  
15 agarose technique, hybrids are seeded in a semi-solid upper layer, over a lower layer containing feeder cells. The colonies from the upper layer may be picked up and eventually transferred to wells.

Antibody-secreting hybrids can be grown in various tissue culture flasks, yielding supernatants with variable concentrations of antibodies. In order to obtain higher  
20 concentrations, hybrids may be transferred into animals to obtain inflammatory ascites. Antibody-containing ascites can be harvested 8-12 days after intraperitoneal injection. The ascites contain a higher concentration of antibodies but include both monoclonals and immunoglobulins from the inflammatory ascites. Antibody purification may then be achieved by, for example, affinity chromatography.

25 Recombinant antibodies are also contemplated, and in particular, single chain antibodies prepared according to Pastan *et al.*, U.S. Patent No. 5,608,039 (hereby incorporated by reference). In particular, humanized antibodies are contemplated. Such antibodies are non-human antibodies in which some or all of the amino acid residues are replaced with the corresponding amino acid residue found in a similar human antibody. Humanization thereby  
30 reduces the antigenic potential of the antibody.

It is not intended that the present invention be limited to specific portions of LAT for the generation of antibodies. However, the cytosolic tail of LAT is a preferred portion for

generating specific antibodies. Such antibodies can be subjected to differential absorption using GST-LAT truncations to enhance specificity for different parts of the molecules.

In another embodiment, antibodies to phosphorylated LAT are contemplated. In such embodiments, phosphorylated expressed LAT protein or synthetically prepared phosphotyrosine containing peptides with the LAT sequence are used. Such antibodies are contemplated as useful for studying (and detecting) the phosphorylation of LAT and are contemplated as antibodies useful for binding activated LAT. Such reagents can be introduced into cells to target specific LAT interactions.

## 2. Antibody Assays

The presence of the LAT polypeptide may be accomplished in a number of ways. A wide range of immunoassay techniques are available as can be seen by reference to U.S. Patent Nos. 4,016,043 and 4,424,279 and 4,018,653 (herein incorporated by reference). This, of course, includes both single-site and two-site, or "sandwich", assays of the non-competitive types, as well as in the traditional competitive binding assays.

Sandwich assays are among the most useful and commonly used assays. A number of variations of the sandwich assay technique exist, and all are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabelled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen secondary complex, a second antibody specific to the antigen, labelled with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of a tertiary complex of antibody-antigen-labelled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, by simple observation of the visible signal, or may be quantitated by comparing with a control sample containing known amounts of hapten. Variations on the forward assay include a simultaneous assay, in which both sample and labelled antibody are added simultaneously to the bound antibody. These techniques are well known to those skilled in the art, including any minor variations as will be readily apparent.

In the typical forward sandwich assay, a first antibody having specificity for LAT or antigenic parts thereof, is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose.

polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other surface suitable for conducting an immunoassay. The binding processes are well-known in the art and generally consist of cross-linking covalently binding or physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase complex and incubated at 25°C (or higher) for a period of time sufficient to allow binding. The incubation period will vary but will generally be in the range of about 1 minute to 2 hours, and more typically 2-40 minutes. Following the incubation period, the antibody subunit solid phase is washed and dried and incubated with a second antibody. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody.

An alternative method involves immobilizing LAT (or a sample containing LAT) and then exposing the immobilized LAT to specific antibody which may or may not be labelled with a reporter molecule. Depending on the amount of LAT and the strength of the reporter molecule signal, LAT may be detectable by direct labelling with the antibody. Alternatively, a second labelled antibody, specific to the first antibody is exposed to the first antibody/LAT complex to form a tertiary complex. The complex is detected by the signal emitted by the reporter molecule.

By "reporter molecule" as used in the present specification, is meant a molecule which, by its chemical nature, provides an analytically identifiable signal which allows the detection of antigen-bound antibody. Detection may be either qualitative or quantitative. The most commonly used reporter molecules in this type of assay are either enzymes, fluorophores, luminescent molecules or radionuclide containing molecules (i.e. radioisotopes).

In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different conjugation techniques exist, which are readily available to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, amongst others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable color change. For example, p-nitrophenyl phosphate is suitable for use with alkaline phosphatase conjugates; for peroxidase conjugates, 1,2-phenylenediamine, 5-aminosalicylic acid, or toluidine are commonly used. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the

chromogenic substrates noted above. In all cases, the enzyme-labelled antibody is added to the first antibody hapten complex, allowed to bind, and then the excess reagent is washed away. A solution containing the appropriate substrate is then added to the tertiary complex of antibody-antigen-antibody. The substrate will react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of hapten which was present in the sample. "Reporter molecule" also extends to use of cell agglutination or inhibition of agglutination such as red blood cells on latex beads, and the like.

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labelled antibody adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic color visually detectable with a light microscope.

Immunofluorescent and EIA techniques are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radionuclide, chemiluminescent or bioluminescent molecules, may also be employed. It will be readily apparent to the skilled technician how to vary the procedure to suit the required purpose.

### 3. LAT Binding Ligands

The present invention contemplates using LAT, LAT fragments, mutant LAT, and mutant LAT fragments to identify LAT Binding Ligands. Exemplary natural intracellular binding targets include protein tyrosine kinases which comprise one or more LAT binding sites and phosphotyrosine peptide fragments thereof, and protein kinases such as activated ZAP-70 and/or Syk tyrosine kinases and fragments of such targets which are capable of LAT-specific binding. Other natural LAT binding targets and fragments thereof are readily identified by screening cells, membranes and cellular extracts and fractions with the disclosed materials and methods and by other methods known in the art. For example, two-hybrid screening using ZAP-70 and/or Syk kinases are used to identify intracellular targets which specifically bind such LAT fragments. Preferred LAT fragments retain the ability to specifically bind at least one ZAP-70 and/or Syk kinase binding sites.

A wide variety of molecular and biochemical methods are available for generating and expressing LAT fragments and ZAP-70 and/or Syk kinase. For example, the ZAP-70 and/or

Syk kinase and LAT or fragments thereof may be obtained by chemical synthesis. expression in bacteria such as *E. coli* and eukaryotes such as yeast or vaccinia or baculovirus-based expression systems, insect cells etc., depending on the size, nature and quantity of the LAT or fragment. The LAT fragments are of length sufficient to provide a novel peptide. As used  
5 herein, such peptides are at least 5, usually at least about 6, more usually at least about 8, most usually at least about 10 amino acids. LAT fragments may be present in a free state or bound to other components such as blocking groups to chemically insulate reactive groups (e.g. amines, carboxyls, etc.) of the peptide, fusion peptides or polypeptides (i.e. the peptide may be present as a portion of a larger polypeptide). etc. The same applies to the binding  
10 ZAP-70 and/or Syk kinase fragments, other LAT binding ligands etc.

The LAT fragments maintain binding affinity of not less than six, preferably not less than four, more preferably not less than two orders of magnitude less than the binding equilibrium constant of a full-length native LAT to the binding target under similar conditions.

15 Particular LAT fragments or deletion mutants are shown to function in a dominant-negative fashion. Overexpression of particular LAT fragments, for e.g. mLAT are also shown to block NFAT and AP-1 transcriptional activation mediated by OKT3 stimulated Jurkat T cells. Such LAT fragments provide therapeutic agents, e.g. when delivered by intracellular immunization--transfection of susceptible cells with nucleic acids encoding such  
20 mutants. The claimed LAT and LAT fragments are isolated, partially pure or pure and are typically recombinantly produced. As used herein, an "isolated" peptide is unaccompanied by at least some of the material with which it is associated in its natural state and constitutes at least about 0.5%, preferably at least about 2%, and more preferably at least about 5% by weight of the total protein (including peptide) in a given sample; a partially pure peptide  
25 constitutes at least about 10%, preferably at least about 30%, and more preferably at least about 60% by weight of the total protein in a given sample; and a pure peptide constitutes at least about 70%, preferably at least about 90%, and more preferably at least about 95% by weight of the total protein in a given sample.

30 The invention provides LAT-specific binding agents, methods of identifying and testing such agents, and their use in diagnosis, therapy and pharmaceutical development. For example, LAT-specific agents are useful in a variety of diagnostic applications, especially where disease or disease prognosis is associated with immune dysfunction resulting from improper expression of LAT or due to the signaling pathways being modified by LAT. In

addition. LAT specific agents are useful in detecting clinical conditions of LAT deficiency. LAT mutations by assessing the levels of LAT expression and/or LAT phosphorylation in clinical samples, by developing assays such as PCR, Immunoblotting etc. Novel LAT-specific binding agents include LAT-specific antibodies; novel nucleic acids with sequence similarity to that of LAT; isolated ZAP-70 and/or Syk kinase binding domains; Grb2 binding domains; other natural intracellular binding agents identified with assays such as one- and two-hybrid screens; non-natural intracellular binding agents identified in screens of chemical libraries, etc.

The invention also provides nucleic acids encoding the subject LAT and LAT fragments thereof. the said nucleic acids may be part of LAT-expression vectors and may be incorporated into recombinant cells for expression and screening, transgenic animals for functional studies, etc. In addition, the invention provides nucleic acids sharing substantial sequence similarity with that of one or more wild-type (WT) LAT nucleic acids. Substantially identical or homologous nucleic acid sequences hybridize to their respective complements under high stringency conditions, for example, at 55°C and hybridization buffer, comprising 50% formamide in 0.9M saline/0.09M sodium citrate (SSC) buffer and remain bound when subject to washing at 55°C with the SSC/formamide buffer. Where the sequences diverge, the differences are preferably silent, i.e. or a nucleotide change providing a redundant codon, or conservative, i.e. a nucleotide change providing a conservative amino acid substitution.

The nucleic acids find a wide variety of applications including use as hybridization probes, PCR primers, therapeutic nucleic acids, etc. for use in detecting the presence of LAT genes and gene transcripts, for detecting or amplifying nucleic acids with substantial sequence similarity such as LAT homologs and structural analogs, and for gene therapy applications (including antisense approaches). Given the subject probes, materials and methods for probing cDNA and genetic libraries and recovering homologs are known in the art. Preferred libraries are derived from human immune cells, especially cDNA libraries from differentiated and activated human lymphoid cells. In one application, the subject nucleic acids find use as hybridization probes for identifying LAT cDNA homologs with substantial sequence similarity. These homologs in turn provide additional LAT and LAT fragments for use in binding assays and therapy as described herein.

Therapeutic LAT nucleic acids are used to modulate, usually reduce, cellular expression or intracellular concentration or availability of active LAT. These nucleic acids are typically antisense: single-stranded sequences comprising complements of the disclosed LAT

nucleic acids. Antisense modulation of LAT expression may employ LAT antisense nucleic acids operably linked to gene regulatory sequences. Cells are transfected with a vector comprising an LAT sequence with a promoter sequence oriented such that transcription of the gene yields an antisense transcript capable of binding to endogenous LAT encoding mRNA.

5 Transcription of the antisense nucleic acid may be constitutive or inducible and the vector may provide for stable extrachromosomal maintenance or integration. Alternatively, single-stranded antisense nucleic acids that bind to genomic DNA or mRNA encoding a LAT or human LAT (hLAT) fragment may be administered to the target cell, in or temporarily isolated from a host, at a concentration that results in a substantial reduction in hLAT

10 expression. For gene therapy involving the transfusion of hLAT transfected cells, administration will depend on a number of variables that are ascertained empirically. For example, the number of cells will vary depending on the stability of the transfused cells. Transfusion media is typically a buffered saline solution or other pharmacologically acceptable solution. Similarly the amount of other administered compositions, e.g. transfected

15 nucleic acid, protein, etc., will depend on the manner of administration, purpose of the therapy, and the like.

The subject nucleic acids are often recombinant, meaning they comprise a sequence joined to a nucleotide other than that which it is joined to on a natural chromosome. An isolated nucleic acid constitutes at least about 0.5%, preferably at least about 2%, and more

20 preferably at least about 5% by weight of total nucleic acid present in a given fraction. A partially pure nucleic acid constitutes at least about 10%, preferably at least about 30%, and more preferably at least about 60% by weight of total nucleic acid present in a given fraction. A pure nucleic acid constitutes at least about 80%, preferably at least about 90%, and more preferably at least about 95% by weight of total nucleic acid present in a given fraction.

25 The invention provides efficient methods of identifying pharmacological agents or drugs which are active at the level of LAT modulatable cellular function, particularly LAT mediated T cell antigen receptor signal transduction. Generally, these screening methods involve assaying for compounds which interfere with hLAT activity such as hLAT-ZAP-70 and/or LAT-Syk kinase binding, LAT-Grb2 binding, etc. The methods are amenable to

30 automated, cost-effective high throughput drug screening and have immediate application in a broad range of domestic and international pharmaceutical and biotechnology drug development programs. Target therapeutic indications are limited only in that the target cellular function (e.g. gene expression) be subject to modulation, usually inhibition, by

disruption of the formation of a complex comprising a hLAT or hLAT fragment and one or more natural hLAT intracellular binding targets. Since a wide variety of genes are subject to ZAP-70 and/or Syk kinase regulated signaling, target indications may include viral, bacterial and fungal infections, metabolic disease, genetic disease, cell growth and regulatory disfunction, such as neoplasia, inflammation, hypersensitivity, etc. Frequently, the target indication is related to either immune dysfunction or selective immune suppression.

A wide variety of assays for binding agents are provided including labelled *in vitro* protein-protein and immunoassays for protein binding or complex formation, cell based assays such as two or three hybrid screens, transient transfection and co-immunoprecipitation assays, etc. For example, three-hybrid screens are used to rapidly examine the effect of transfected nucleic acids, which may, for example, encode combinatorial peptide libraries or antisense molecules, on the intracellular binding of hLAT or hLAT fragments to intracellular hLAT targets. Convenient reagents for such assays (e.g. GST fusion partners) are available in the art.

hLAT or hLAT fragments used in the methods are usually added in an isolated, partially pure or pure form and are typically recombinantly produced. The hLAT or fragment may be part of a fusion product with another peptide or polypeptide, e.g. a polypeptide that is capable of providing or enhancing protein-protein binding, sequence-specific nucleic acid binding or stability under assay conditions (e.g. a tag for detection or anchoring). The assay mixtures comprise at least a portion of a natural intracellular hLAT binding target such as ZAP-70 and/or Syk kinase subunit domain or Grb2 binding domains. The assay mixture also comprises a candidate pharmacological agent. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e. at zero concentration or below the limits of assay detection. Candidate agents encompass numerous chemical classes, though typically they are organic compounds; preferably small organic compounds. Small organic compounds have a molecular weight of more than 50 yet less than about 2,500, preferably less than about 1000, more preferably, less than about 500. Candidate agents comprise functional chemical groups necessary for structural interactions with proteins and/or DNA, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups, more preferably at least three. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the aforementioned functional groups.



Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof, and the like. Where the agent is or is encoded by a transfected nucleic acid, said nucleic acid is typically DNA or RNA. Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural and synthetically produced libraries and compounds are readily modified through conventional chemical, physical, and biochemical means. In addition, known pharmacological agents may be subject to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc., to produce structural analogs. A variety of other reagents may also be included in the mixture. These include reagents like salts, buffers, neutral proteins, e.g. albumin, detergents, etc. which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions, etc. Also, reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, antimicrobial agents, etc. may be used. The resultant mixture is incubated under conditions whereby, but for the presence of the candidate pharmacological agent, the LAT specifically binds the cellular binding target, portion or analog. The mixture components can be added in any order that provides for the requisite bindings. Incubations may be performed at any temperature which facilitates optimal binding, typically between 4 and 40°C, more commonly between 15 and 40°C. Incubation periods are likewise selected for optimal binding but also minimized to facilitate rapid, high-throughput screening, and are typically between 0.1 and 10 hours, preferably less than 5 hours, more preferably less than 2 hours. After incubation, the presence or absence of specific binding between the hLAT and one or more binding targets is detected by any convenient way. For cell-free binding type assays, a separation step is often used to separate bound from unbound components. The separation step may be accomplished in a variety of ways. Conveniently, at least one of the components is immobilized on a solid substrate which may be any solid from which the unbound components may be conveniently separated. The solid substrate may be made of a wide variety of materials and in a wide variety of shapes, e.g. microtiter plate, microbead, dipstick, resin particle, etc. The substrate is chosen to maximize signal to noise ratios, primarily to minimize background binding, for ease of washing and cost. Separation

may be effected for example, by removing a bead or dipstick from a reservoir, emptying or diluting reservoir such as a microtiter plate well, rinsing a bead (e.g. beads with iron cores may be readily isolated and washed using magnets), particle, chromatographic column or filter with a wash solution or solvent. Typically, the separation step will include an extended  
5 rinse or wash or a plurality of rinses or washes. For example, where the solid substrate is a microtiter plate, the wells may be washed several times with a washing solution, which typically includes those components of the incubation mixture that do not participate in specific binding such as salts, buffer, detergent, nonspecific protein, etc. may exploit a polypeptide specific binding reagent such as an antibody or receptor specific to a ligand of  
10 the polypeptide. Detection may be effected in any convenient way. For cell based assays such as one, two, and three hybrid screens, the transcript resulting from hLAT-target binding usually encodes a directly or indirectly detectable product (e.g. galactosidase activity, luciferase activity, etc.). For cell-free binding assays, one of the components usually comprises or is coupled to a label. A wide variety of labels may be employed, essentially any  
15 label that provides for detection of bound protein. The label may provide for direct detection as radioactivity, luminescence, optical or electron density, etc. or indirect detection such as an epitope tag, an enzyme, etc. The label may be appended to the protein e.g. a phosphate group comprising a radioactive isotope of phosphorous, or incorporated into the protein structure, e.g. a methionine residue comprising a radioactive isotope of sulfur. A variety of methods  
20 may be used to detect the label depending on the nature of the label and other assay components. For example, the label may be detected bound to the solid substrate or a portion of the bound complex containing the label may be separated from the solid substrate, and thereafter the label detected. Labels may be directly detected through optical or electron density, radiative emissions, nonradiative energy transfers, etc. or indirectly detected with  
25 antibody conjugates, etc. For example, in the case of radioactive labels, emissions may be detected directly, e.g. with particle counters or indirectly, e.g. with scintillation cocktails and counters. The methods are particularly suited to automated high throughput drug screening. Candidate agents shown to inhibit hLAT-target binding provide valuable reagents to the pharmaceutical industries for animal and human trials. For example, LAT fragments capable  
30 of blocking T cell activation, find use in treating disease associated with undesirable cell growth, differentiation, particularly immune cell differentiation, and hypersensitivity/allergy. In addition, they find use in the inhibition of autoreactive T cells (autoimmunity), inhibition of production of deleterious T cell products (such as cytokines, lymphokines) and/or T cell

activities (such as cytotoxicity, etc). Also, the invention finds use in treating diseases associated with cell movement, particularly as changes in motility are important in growth and differentiation. Thus, it could be relevant in the treatment of cancer, immunological diseases, autoimmune diseases, graft rejection and others. For therapeutic uses, the compositions and agents disclosed herein may be administered by any convenient way, preferably parenterally, conveniently in a physiologically acceptable carrier, e.g., phosphate buffered saline, saline, deionized water, or the like. Typically, the compositions are added to a retained physiological fluid such as blood or synovial fluid. Generally, the amount administered will be empirically determined, typically in the range of about 10 to 1000  $\mu\text{g/kg}$  of the recipient. For peptide agents, the concentration will generally be in the range of about 100 to 500  $\mu\text{g/ml}$  in the dose administered. Other additives may be included, such as stabilizers, bactericides, etc. These additives will be present in conventional amounts.

#### DETAILED DESCRIPTION OF SPECIFIC EMBODIMENTS

Generally, the nomenclature used hereafter and the laboratory procedures in cell culture, molecular genetics, and nucleic acid chemistry and hybridization described below are those well known and commonly employed in the art. Standard techniques are used for recombinant nucleic acid methods, polynucleotide synthesis, and microbial culture and transformation (e.g., electroporation, lipofection). Generally enzymatic reactions and purification steps are performed according to the manufacturer's specifications. The techniques and procedures are generally performed according to conventional methods in the art and various general references (see, generally, Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2d ed. (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., and *Current Protocols in Molecular Biology* (1996) John Wiley and Sons, Inc., N.Y.) which is incorporated herein by reference) which are provided throughout this document. All the information contained therein is incorporated herein by reference.

Oligonucleotides can be synthesized on an Applied BioSystems oligonucleotide synthesizer [for details see Sinha *et al.*, *Nucleic Acids Res.* 12:4539 (1984)], according to specifications provided by the manufacturer. Complementary oligonucleotides are annealed by heating them to 90°C in a solution of 10 mM Tris-HCl buffer (pH 8.0) containing NaCl (200 mM) and then allowing them to cool slowly to room temperature. For binding and turnover assays, duplex DNA is purified from native polyacrylamide (15% w/v) gels. The band corresponding to double-stranded DNA is excised and soaked overnight in 0.30 M sodium

acetate buffer (pH 5.0) containing EDTA (1 mM). After soaking, the supernatant is extracted with phenol/chloroform (1/1 v/v) and precipitated with ethanol. DNA substrates are radiolabeled on their 5'-OH group by treatment with [ $\gamma$ - $^{32}$ P]ATP and T4 polynucleotide kinase. Salts and unincorporated nucleotides are removed by chromatography on Sephadex G  
5 columns.

Assays for detecting the ability of agents to inhibit or enhance LAT binding provide for facile high-throughput screening of agent banks (e.g., compound libraries, peptide libraries, and the like) to identify antagonists or agonists. Such ZAP-70 and/or syk tyrosine kinase signaling pathway antagonists and agonists may be further developed as potential  
10 therapeutics and diagnostic or prognostic tools for diverse types of infectious diseases, autoimmune diseases and hereditary diseases, and cancer.

#### A. Drug Screening Assays for Agonists and Antagonists

In some embodiments, LAT (and in particular, fragments of LAT) are useful in drug  
15 screening assays designed to identify drugs that interfere with the specific binding of ZAP-70 and/or Syk kinases with its substrate as well as ZAP-70 and/or Syk kinase activity, and thereby block the activation of downstream signaling molecules.

In preferred embodiments, the invention provides an isolated LAT polypeptide, or a fragment thereof, having Grb2-specific binding affinity. The invention provides nucleic acids  
20 encoding the LAT polypeptide and LAT fragments as part of expression vectors for introduction into cells. The invention provides methods of identifying intracellular molecules which interact with LAT or LAT fragments, as well as exogenous agents (*i.e.* drugs) which disrupt or enhance the binding of LAT and/or fragments thereof to such intracellular targets.

The claimed polypeptide LAT and LAT fragments find particular use in screening  
25 assays for agents or lead compounds for agents useful in the diagnosis, prognosis or treatment of disease, particularly disease associated with undesirable cell growth, cell movement, differentiation and cell proliferation. One such assay involves forming mixtures of 1) LAT (or fragments thereof) and 2) an intracellular LAT-binding ligand, in the presence or absence of 3) a prospective drug candidate. The mixtures are made under conditions that permit the  
30 binding of the intracellular LAT-binding ligand to LAT (or fragments thereof) and the mixtures are then analyzed for the presence of such binding. A difference in such binding in the presence of such a drug candidate indicates that the agent is capable of modulating the binding of LAT (or fragments thereof) to an intracellular LAT-binding ligand. The assays of

the present invention provide for facile high-throughput screening of compounds suspected to be able to inhibit such binding (e.g., compound libraries, peptide libraries, and the like) to identify potential drug candidates.

The compositions and methods of the present invention can be used to identify and screen for drugs that block the following LAT interactions (also see detailed description of specific embodiments):

- 1) Development of drugs that prevent ZAP-70 and/or other kinases from interacting and phosphorylating LAT
- 2) Development of drugs that prevent LAT from interacting with ZAP-70 or from getting phosphorylated.
- 3) Development of drugs that prevent binding of phosphorylated LAT to ligands such as Grb2 etc.
- 4) Development of drugs that prevent binding of Grb2 (or other ligands) to phosphorylated LAT.

An example of a drug screening assay, that can be employed to test for potential drug candidates that inhibit the specific binding of Grb2 with LAT, is given below. The suspect drug candidates are tested whether they block the binding of Grb2 to the full length LAT, or preferably to Grb2 fused to glutathione-S-transferase (GST). GST-Grb2 or GST-Grb2 is immobilized on glutathione-agarose beads and incubated for 1 hour with the indicated amount or different doses of the drug candidate. In parallel assays, the mutant LAT or  $\alpha$ LAT antibodies are used as controls. Lysates from Jurkat T cells treated for 2min with or without C305 are incubated with the immobilized GST-Grb2 in the presence or absence of the drug candidate (or in parallel assays with the mutant LAT). Bound proteins are eluted with sodium dodecyl sulfate-containing buffer, separated by SDS-PAGE and immunoblotted with  $\alpha$ LAT. GST-Grb2 binds to LAT in C305 stimulated Jurkats, but not unstimulated cells. Alternatively, one can screen for drugs that block Grb2 binding to GST-LAT. Treatment with mutant LAT (mLAT) or  $\alpha$ LAT antibodies inhibits the binding of GST-Grb2 to LAT in a dose-dependent fashion.

In other embodiments, drug screening assays that can be employed for testing potential drug candidates (for agonists), are those that overcome the inhibition of transcriptional activation of NF-AT (involved in TCR-mediated transcriptional events). Details of these assays are given below. Jurkat TAg cells, can be stably transfected with mutant LAT(Y171/191F), (in parallel assays with the wild-type or vector only), together with a

reporter construct for secreted alkaline phosphatase (SEAP) driven by the NF-AT response elements. The suspect drug candidates are tested whether they overcome the inhibition of the transcriptional activation of NF-AT in OKT3 stimulated mLAT transfected Jurkat cells. (For more details, see methodology and example 7 in the experimental section). Identification of agonists are particularly useful in treatment of clinical conditions of LAT deficiency.

## B. Structural and Functional Investigation of LAT.

In other embodiments, signaling molecules that bind to LAT can be identified. Based on structural and functional features of LAT (See below and experimental Section). LAT may serve as an adapter molecule that recruits multiple signaling molecules to Ligand-LAT complexes. It is believed that one or more of these proteins may link these complexes to cellular functions. Signaling proteins are predicted either to bind to pTyr within LAT in response to ligand or to bind constitutively to LAT. Identification of LAT binding partners also would provide insight into T cell, NK cell and mast cell signaling pathways. The techniques for studying the interaction of the protein(s) of interest with LAT and analyzing their role in cellular functions are as described below and in the experimental section.

### i. Structure-function analysis of LAT

In some embodiments, LAT can be subjected to mutagenesis followed by analysis of function, i.e., to understand how does LAT regulate the following and how do mutations of LAT affect the following :

A. **Functional assays:** -all can be performed in T cells (Jurkat, normal T cells from humans or mice), mast cells, NK cells. Alternatively, non-lymphocyte cells can be studied by transient or stable expression of LAT or LAT mutants. Additionally one can express necessary kinases, receptors, accessory molecule and relevant indicator molecules. Studies can be performed *in vitro*, in intact cells or in animal models.

#### i). **Protein kinase assays**

a. protein tyrosine kinases-Lck, Fyn, ZAP-70, Syk, Itk  
b. protein serine kinases-Raf, MEK (all), ERK (all), PAK, MEKK (all), JNKK (all), JNK (all) and all related identified and, as of now, unidentified members of these protein serine/threonine kinase families. Protein kinase C (all family members). One can identify novel kinases regulated directly or indirectly by LAT.

ii). **Calcium assays**-one can analyze elevations from extra-and intra-cellular sites

iii). *Assays of small G proteins*- (including, but not limited to Ras, Rac, Rho, cdc42, Rap) and regulators of small G protein function (including, but not limited to SOS, GAP, Vav)

iv). *Assays of protein binding* -including, but not limited to those molecules shown to be directly or indirectly bound to LAT (Grb2, Grap, phospholipase C, PI-3 kinase, Vav, SLP-76, Cbl)

v). *Assays of transcriptional regulation*- including, but not limited to NFAT, AP-1, Fos, Jun

vi). *Localization assays using indirect immunofluorescence and GFP chimeras*.  
Assay by light, confocal, imaging microscopy

vii). *Assays of T cell cellular function*- including, but not limited to cell shape, cell size, cell motility, cell migration, cell adhesion

viii). *Assays of tyrosine and serine/threonine phosphorylation of LAT*- By using mass spectrometry, chromatography and antibodies specific for all regions of LAT

ix). *Generate panels of anti-LAT antibodies*- antibodies that bind phosphorylated and non-phosphorylated regions of LAT.

## **B. Mutagenesis of LAT**

In preferred embodiments, Structure-function analysis of LAT can be performed by mutagenesis of: i) Tyrosine residues, singly and in combinations; ii) Truncations from carboxy end and truncations of amino terminal end of the cytosolic domain (the cytosolic domain is defined approximately by amino acids 28 to 233) followed by re-generation of in-frame sequence; point mutations as indicated following truncations; iii) Deletion of putative transmembrane domain (the transmembrane domain is defined approximately by residues 5 to 27) and mutagenesis of residues in putative transmembrane domain; iv) Mutations of charged residues and cysteine residues near putative transmembrane domain.

## **ii. Analysis of LAT interactions**

In other embodiments, analysis of LAT interactions with other signaling proteins can be performed as given below.

A. One can study the interactions of LAT with the signaling molecules as described in the above section and in the experimental section. Also, one can map sites of interactions as stated above (mutagenesis, region-specific antibodies, etc.).

B. One can determine new and additional interactions with additional intracellular molecules. by Protein purification, such as Affinity purification using LATprotein. mRNA or cDNA as a probe. By employing two and three hybrid analysis to identify interacting molecules.

5 C. One can determine interactions with integral membrane proteins including. but not limited to any component of the T cell, B cell, mast cell or NK cell antigen or immunoglobulin receptors, including, but not limited to accessory or co-receptor molecules such as CD4, CD8, CD28.

10 **iii. Analysis of the role of LAT in T cell, mast cell and NK cell function**

In some embodiments, the function of LAT can be ascertained in various cells as given below.

A. In T cells, one can study T cell activation, T cell apoptosis, T cell anergy, T cell development and differentiation, T cell helper and killer function, T cell lymphocyte/cytokine production. In addition, one can generate transgenic animals overexpressing wild-type or mutant LAT; generation of LAT knock-outs. Also, one can generation LAT deficient T cell line as recipient for mutant forms of LAT.

B. In Mast cells, one can study granule release, cytokine production, second messenger production, gene transcription

20 C. In NK cells one can study various functions, such as cytotoxicity, cytokine and second messenger production, gene transcription

**iv. Structural analysis of LAT**

In other embodiments, structural analysis of LAT can be performed as given below.

25 A. Mutation of LAT to generate cytosolic fragment followed by analysis of structure by NMR and/or crystallography.

B. Comparison of structure of mutant forms of LAT. Comparison of structure of non-phosphorylated and phosphorylated forms of LAT.

C. Modeling small molecule interactions with structural models of LAT.

30 **iv. Inhibitors of LAT function**

In preferred embodiments, inhibitors of LAT function can be identified and/or generated as given below.



A. Generation of antibodies that bind LAT and block function. including monoclonal, single chain antibodies, recombinant antibodies

B. Identify small molecules that bind LAT, by screening libraries of organics for binders, peptide binders, molecular design

C. Generation of anti-sense molecules to decrease LAT expression

D. Assay as above for function, interaction, localization, etc.

vi. **LAT-mediated immunodeficiency**

In preferred embodiments, the compositions and methods of the present invention can be employed for developing diagnostic assays

A. Screen various patients with immunodeficiencies to find patients lacking LAT or with mutated LAT. Define clinical condition(s) characterized by LAT deficiency

B. Establish cell lines from such patients. Reconstitute with type. Define mutations in LAT; reexpress mutant form (patient form) in LAT deficient cell line. Use naturally occurring mutants for structure-function analysis.

C. Develop screening test for LAT deficiency or mutation.

D. Develop genetic therapy for LAT deficiency; bone marrow or T cell reconstitution.

## EXPERIMENTAL

The following examples serve to illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof. In the experimental disclosure which follows, the following methodology apply:

**Antibodies, Immunoprecipitation and Western Blotting:** Antibodies used in the experiments were rabbit polyclonal anti-Grb2, anti-Grb2 cross-linked to agarose, anti-Cbl (C-15), and anti-Sam68 from Santa Cruz Biotechnology. Monoclonal anti-PLC- $\gamma$ 1, anti-phosphotyrosine (4G10) and anti-Vav, rabbit anti-p85 polyclonal were from Upstate Biotechnology. Others were anti-Flag M2 (Kodak), and anti-phosphotyrosine antibody (PY20, Transduction Laboratory). PY20 was chemically cross-linked to protein A-sepharose using dimethyl pimelimidate (DMP). GST fusion proteins were crosslinked to glutathione-sepharose with 1-Ethyl-3-(3-Dimethylaminopropyl) carbodiimide (EDC, Pierce). Anti-LAT antibodies were generated against a GST fusion protein containing the cytosolic portion of LAT (aa

31-233). The following antibodies were used: rabbit polyclonal anti-LAT antiserum (Zhang, *et al.*, LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation. *Cell* 92:83-92, 1998), anti-ZAP-70 (Wange, *et al.*, Activating and inhibitory mutations in adjacent tyrosines in the kinase domain of ZAP-70, *J. Biol. Chem.* 270:18730-18733, 1995),  
5 anti-Lck (Samelson *et al.*, Association of the fyn protein-tyrosine kinase with the T-cell antigen receptor, *Proc. Natl. Acad. Sci. USA* 87:4358-4362, 1990), anti-CD3 $\epsilon$  (OKT3, Kung *et al.*, Monoclonal antibodies defining distinctive human T cell surface antigens. *Science* 206:347-349, 1979), monoclonal anti-myc (9E10) and monoclonal anti-PLC $\gamma$ 1 (a gift from Dr. S. G. Rhee, NHLBI, NIH).

10 Jurkat T cells (ATCC) were removed from culture, washed and resuspended at  $10^8$  cells/ml in RPMI. Cells were either stimulated with OKT3 ascites (1:100) (ATCC) or C305 (1:50) for 2 minutes or left untreated and lysed in 1% Brij lysis buffer (Wange *et al.*, Activating and inhibitory mutations in adjacent tyrosines in the kinase domain of ZAP-70. *J. Biol. Chem.* 270, 18730-18733, 1995).

15 For treatment with calf intestine alkaline phosphatase (CIP), immunoprecipitates from  $10^7$  cells were incubated with 10 units of CIP (Boehringer) in 40  $\mu$ l CIP reaction buffer for 1 hour.


**Purification of LAT:**  $10^{11}$  Jurkat E6.1 cells were stimulated for 1.5 min at  $10^8$ /ml in RPMI-1640 with OKT3 ascites (1:100) at 37 °C. Immediately after stimulation, cells were  
20 washed with cold RPMI quickly and membrane fractions were prepared by Dounce homogenization and centrifugation. Membrane fractions were then extracted with 1% Brij lysis buffer, and insoluble material was removed by centrifugation. Membrane protein extracts were heated for 10 min at 75°C, followed by centrifugation to remove insoluble material. This supernatant was loaded onto a PY20 column, which was washed sequentially  
25 with lysis buffer containing 1% Brij 97 (25mM Tris-HCl pH 7.6, 150mM NaCl, 5mM EDTA, 1mM sodium orthovanadate, 10  $\mu$ g/ml each of leupeptin, aprotinin, p-nitrophenyl-p-guanidinobenzoate), 1% CHAPS (3-[(3-cholamidopropyl)-dimethylammonio]-l-propanesulfonate), and 0.1% CHAPS. Phospho-tyrosine containing proteins were eluted with  
30 0.1 % CHAPS buffer containing 10mM phenyl phosphate. Eluted proteins were concentrated and subjected to SDS-PAGE. The band for p36-38 was excised and subjected to *in gel* S-carboxyamidomethylation followed by tryptic digestion as described, without the addition of 0.02% Tween. The resulting peptide mixture was separated by microbore HPLC using a Zorbax C18 1.0mm by 150 mm reverse-phase column on a Hewlett-Packard 1090

HPLC/1040 diode array detector. Optimum fractions were chosen based on differential UV absorbance at 205nm, 277nm and 292nm, peak symmetry and resolution: then further screened for length and homogeneity by matrix-assisted laser desorption time-of-flight spectrometry (MALDI-MS) on a Finnigan Lasermat 2000 (Hemel England). Tryptic peptides were submitted to automated Edman degradation on an Applied Biosystems Procise 494 or 477A protein sequencer (Foster City, CA).

**cDNA cloning and Northern blotting:** A YT lambda ZAP cDNA library (10<sup>6</sup> plaques) was screened with a random primed probe made from the *Eco* RI/*Pst* I fragment of an EST clone (Genbank #w74254). 20 overlapping phage clones were isolated. The cDNAs were excised from lambda phages with the rapid excision kit from Stratagene. For Northern analysis, blots with poly(A)<sup>+</sup> RNA from different human tissues (ClonTech) were probed with a radiolabeled fragment of LAT cDNA or human  $\beta$ -actin cDNA under high stringency conditions according to the manufacturer's instruction.

**Construction of LAT in expression vector and mutagenesis:** The myc tag was fused to the C-terminus of LAT by replacing the *Pst* I/*Hind* III fragment of LAT in pBluescript (SK-) (Stratagene) with a double stranded oligonucleotide fragment encoding the myc tag sequence (SMEQKLISEEDLN). Flag-tagged LAT was constructed by replacing the *Pst* I/*Cla* I fragment of LAT in pBluescript with a double stranded oligonucleotide fragment encoding the FLAG sequence (DYKDDDDK). The tagged LAT cDNAs were cloned into the pcDNA3 (Invitrogen) or pEF/BOS vector (See Schematic B, and Mizushima and Nagata *et al.*, *Nuc. Acid Res.*, 18: 5322, 1990).

As shown in Schematic B, pEF-BOS carries the SV40 replication origin (311 bp of *Eco*RII G fragment), the promoter region of human EF-1 $\alpha$  chromosomal gene (1.2 kb), the stuffer fragment (450 bp) from CDM8 vector (See Seed, B. *Nature* 329:840-842, 1987), and poly(A) adenylation signal from human EF-1 $\alpha$  chromosomal gene (1.2 kb), the stuffer fragment (450 bp) from CDM8 vector and poly(A) adenylation signal from human G-CSF cDNA (700 bp *Eco*811~*Eco*RI DNA fragment) (Nagata S *et al.*, *Nature* 319: 415-418, 1986) *avin Hind*III-*Eco*RI site of pUC119. The promoter region of EF-1 $\alpha$  gene is from nucleotide position 373 to 1561 (Uetsuki *et al.*, *J. Biol. Chem.*, 264:5791-5798, 1989), which includes 203 bp 5' flanking region, 33 bp first exon, 943 bp first intron and 10bp of the part of the second exon located at 20 bp upstream of the ATG codon. The size of the pEF-BOS is 5.8 kb. and the cDNA to be expressed can be inserted at *Bst*XI site using *Bst*XI adapter, or *Xba*I site using *Xba*I linker. Mutagenesis of Y171 to F and Y191 to F was done by PCR.



The structure of pEF-BOS. The boxes indicate SV40 origin, human EF-1 $\alpha$  promoter region, stuffer from CDM8 and poly(A) adenylation site, respectively. The slashed areas in the EF-1 $\alpha$  promoter region are first exon and the part of second exon, respectively. The lines flanking boxes are the sequence of pUC119. Major recognition sites for restriction enzymes are shown.

## Schematic B

**Stable and transient transfections:** Transfection of 293T cells was by the calcium phosphate method. Cells were harvested 36 hours after transfection. For transient transfection of Jurkat/Tag cells,  $2 \times 10^7$  cells in 0.4 ml RPMI-1640, 25 mM Hepes, 2 mM Glutamine were incubated with pSX-NFAT/SEAP or pSX-AP-1/SEAP with 20  $\mu$ g of pEF/LAT( $\omega$ 1).  
5 pEF/LAT(Y171/191F) or pEF/BOS vector DNA as control. 24 hours after transfection, transfected cells were stimulated with OKT3 ascites (1:1000) or PMA (10 ng/ml) plus ionomycin (Sigma) (1.5  $\mu$ M). SEAP assay was done as described (Spencer *et al.*, Controlling signal transduction with synthetic ligands. *Science* 262, 1019-1024, 1993; Berger *et al.*, Secreted placental alkaline phosphatase: a powerful new quantitative indicator of gene  
10 expression in eukaryotic cells, *Gene* 66:1-10, 1988). For stable transfection of Jurkat E6.1,  $10^7$  cells in 0.4 ml RPMI were electroporated using the same conditions as above. Stable transfectants were selected in the presence of 1.5 mg/ml G418 (Gibco/BRL).

**Immunofluorescence Staining and Confocal Microscopy:** HeLa cells transfected with the cDNA for LAT were grown on sterile glass coverslips overnight prior to antibody  
15 staining. Jurkat T cells were stained in suspension, and the cells were mounted onto coverslips immediately prior to analysis. Immunofluorescence staining were performed as described (Sloan-Lancaster *et al.*, Regulation of ZAP-70 intracellular localization: visualization with the green fluorescent protein. *J. Exp. Med.* 186, 1713-1724, 1997) with anti-LAT at 1:500 dilution for T cells or 1:1000 dilution for HeLa cells (ATCC) . or  
20 monoclonal anti-Lck (Santa Cruz Biotech, CA) at 1:500 dilution.

**Ligands of LAT:** The nucleic acid and amino acid sequences for ZAP-70 (GENBANK ACCESSION No. L05148), Syk Kinase (GENBANK ACCESSION No. Z29630), Grb2 (GENBANK ACCESSION No. M96995), Vav (GENBANK ACCESSION No.X16316) and Cbl (GENBANK ACCESSION No.X57110) are available.

25 **Purification of GEM Fractions and Subcellular Fractionation:**  $5 \times 10^7$  cells were lysed on ice in 1 ml 1% Triton in MNE buffer ( 25 mM MES pH 6.5, 150 mM NaCl, 5 mM EDTA). dounced 10 times, and mixed with 1 ml 80% sucrose made with MNE buffer. After transfer of the lysate to the centrifuge tube, 2 ml 30% sucrose in MNE buffer was overlaid, then 1 ml 5% sucrose in MNE was overlaid. After centrifugation for 16-18 hours at  
30 200,000g in a Beckman SW55Ti, 0.4 ml gradient fractions were collected from the top of the gradient. For purification of GEMs from OKT3 stimulated cells, cells were spun down quickly after stimulation and then lysed in 1 ml 1% Triton in 25 mM Tris-Cl pH 7.6, 150 mM NaCl, 5 mM EDTA, 30 mM pyrophosphate, 10 mM glycerol phosphate and 1 mM

sodium orthovanadate. Lysates were prepared for sucrose gradient ultracentrifugation as above.

**Mutagenesis and Subcloning:** Cysteine to alanine mutations (position 26 and 29 ) of LAT cloned into the pCEFL expression vector (a gift from Dr. S. Gutkind, NIDR, NIH) were made with the Stratagene Quickchange kit. The LAT transmembrane domain deletion mutant (residues 1-22) was made by cloning a double-stranded linker annealed with two oligonucleotides (AATTCGCCGCCATGGCACTGTGTG and TGCACACACAGTGCCATGGCGGCG), *Apa* I/*Xba* I fragment from pEF/LAT-myc into pCEFL *Eco* RI/*Xba* I sites.

**Transfection, Labeling of Jurkat Cells and 293T Cells, and Immunofluorescence:** Transient and stable transfection of Jurkat cells were performed as described (Zhang, *et al.*, LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation, *Cell* 92:83-92, 1998). For labeling with [<sup>3</sup>H] palmitate, 2X10<sup>7</sup> Jurkat cells were removed from culture and resuspended in 1 ml RPMI 1640 containing 5% dialyzed fetal calf serum (FCS), 5 mM sodium pyruvate and 0.5 mCi [<sup>3</sup>H]-palmitate for 3 hours. 293T cells were transfected with LAT constructs using the calcium phosphate method in 6 well plates. 24 hours after transfection, 293T cells were labeled with 0.5 mCi [<sup>3</sup>H]-palmitate in 1 ml DMEM containing 5% dialyzed FCS and 5 mM sodium pyruvate, for 3 hours. The gels were fixed, treated with Enlightning (Dupont) for 30 min, dried, and exposed to film for 3-4 weeks. Immunofluorescence staining and confocal microscopy were done with 293T cells transfected with LAT constructs as described (Zhang, *et al.*, LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation, *Cell* 92:83-92, 1998).

**Normal Tissues and Cell Populations:** Paraffin embedded normal lymphoid tissues included lymph nodes showing various forms of reactive changes (10 cases), thymuses obtained during cardiac surgery or surrounding thymomas (3), spleens removed after trauma or because of immune thrombocytopenia (4), bone marrows (6), and small intestine (2). In addition, hematopoietic tissues from three embryos aged 11-12 weeks of gestation were analyzed. Freshly frozen samples of reactive lymph nodes (3), spleen (1), and thymus (1) were also used.

Peripheral blood mononuclear cells (PBMC) were isolated from heparinized blood after Ficoll-Hypaque gradient centrifugation and depleted of plastic adherent cells. For purification of polyclonal natural killer (NK) or T cell populations, PBMC were incubated with anti-CD3 monoclonal antibody (JT3A, gift by Dr. A. Moretta, University of Genova.

Italy) for 30 minutes at 4°C. followed by treatment with goat-anti mouse-coated dynabeads (Dynal. Oslo. Norway) for 30 minutes at 4°C. The resulting CD3 negative lymphocyte populations, containing approximately 1% CD3+ cells, 20-30% HLADR+ cells and 70-80% CD16+CD56+ cells, were cultured in rIL-2 (Cetus Corp. Emeryville. CA). In order to obtain polyclonally activated T cell-enriched lymphocyte populations, PBMC were stimulated with 0.1% (vol/vol) PHA (Gipco Ltd. Paisley. Scotland) for 24 hours and then cultured in rIL-2.

**Neoplastic Tissues:** Two-hundred and sixty-four cases of nodal and extranodal hematology neoplasms were gathered from different institutions: all neoplasms had been previously characterized immunophenotypically on paraffin sections. and in many cases on frozen sections as well. All lymphomas were classified according to the International Lymphoma Study Group Classification, and included all major subtypes of Hodgkin's and Non-Hodgkin's lymphomas.

**Fixatives and Tissue and Cell Processing:** Tissue samples had been fixed in various fixatives. including buffered formalin, B5, Bouin, and Hollande. and embedded in paraffin. Bone marrow biopsies were fixed in B5 for 3 hours and decalcified in 0.1 M EDTA disodium salt aqueous solution for 2-8 hours. Fresh tissues were immediately frozen after biopsy in liquid nitrogen and stored at -800C until used.

CD3+CD16-CD56- T cell and CD3-CD16+CD56+ NK cell populations were washed three times in 0.9% NaCl solution, resuspended at a concentration of  $5 \times 10^6$  cells/ml and utilized for cytopsin preparations (100  $\mu$ l/each slide). Slides were air dried for 24 hours. then fixed in -20°C absolute ethanol for 30 minutes, dried and used for immunocytochemical staining.

**Immunostaining of Fixed Sections:** Details on the production and characterization of the rabbit anti-LAT antibody are reported elsewhere (Zhang, *et al.*. LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation, *Cell* 92:83-92, 1998). The immunostaining for LAT was performed on paraffin sections after antigen retrieval in microwave (3 boiling cycles, 5 minutes each at 750 watts power. with an interval of 1 minute between cycles) in citrate buffer, pH 6.0; the polyclonal antibody anti-LAT was applied at a dilution of 1:800 in TRIS-HCl buffer, pH 7.2-7.4, for 45 minutes, and was followed by biotinylated anti-rabbit antibody (30 minutes) and peroxidase-conjugated streptavidin-biotin complex (30 minutes) (Bio-S.P.A., Milan, Italy). On cryostat sections. the procedure was similar, but the microwave heating was avoided. Sections were air dried for 18 hours. and

LAT was applied at the dilution of 1:100. On cytopins, the sample was subjected to microwave heating once.

In all cases of T-cell lymphomas and anaplastic large cell lymphomas (ALCL), serial paraffin sections were also stained with a polyclonal antibody anti-CD3 (Dako, Milan) (1:200; microwave antigen retrieval in 1 mM EDTA buffer, pH 8.0, 2x5' cycles). Furthermore, the cases of ALCL were also evaluated for their reactivity with the monoclonal antibody ALK1 (Dako) (1:10; microwave antigen retrieval in citrate buffer, 3x5' cycles), which recognizes a formalin-resistant epitope in the nucleophosmin-anaplastic lymphoma kinase (NPM-ALK) chimeric protein. Both CD3 and ALK1 immunostaining were performed using the same indirect immunoperoxidase technique adopted for LAT. Thymuses from embryos were also stained with polyclonal antibodies anti-CD3 and anti-TdT (Dako) (1:200; microwave antigen retrieval in 1 mM EDTA buffer, pH 8.0, 3x5' cycles, overnight incubation of the primary antibody).

Analysis of the distribution of LAT on PMBC was performed using two-colour fluorescence cytofluorometric analysis (FACS) (Ortho Cytoron Absolute) as previously described [135]. For FACS analysis of LAT in combination with monoclonal antibodies JT3A (IgG2a, anti-CD3), KD1 (IgG2a, anti-CD16) and C218 (IgG1, anti-CD56) (all antibodies provided by Dr. A. Moretta), membranes were permeabilized with 0.2% saponin (Sigma) in phosphate buffered saline (PBS) (pH 7.6) for 5 minutes, after fixation with 4% paraformaldehyde in PBS. As second reagents, fluorescein isothiocyanate (FITC)-conjugated swine anti-rabbit antibody (Dako) and phycoerythrin (PE)-conjugated isotype specific goat anti-mouse antibodies (Southern Biotechnology Associates, Birmingham, AL) were used.

### EXAMPLE 1

In this example, the purification process of LAT is described. Stimulation of the TCR on the Jurkat human T cell line by crosslinking with either C305 (anti-TCR $\beta$ , See Fig.1A) or OKT3 (anti-CD3 $\epsilon$ , not shown) monoclonal antibodies induced the tyrosine phosphorylation of multiple proteins, most prominently, p36-38. To define the role of p36-38 in T cell signaling, the protein was purified from activated Jurkat cells. Since, p36-38 appears to be membrane-associated, membrane fractions were prepared from OKT3 stimulated Jurkat cells and extracted with Brij97 detergent. The extracted membrane proteins were then heat-treated, which induced precipitation of about 2/3 of the protein, but not of p36-38. Anti-phosphotyrosine antibodies were used for affinity purification of



phosphotyrosine-containing proteins. These were then specifically eluted with phenyl phosphate, concentrated, and resolved on SDS-PAGE. The p36-38 band was excised, digested *in gel* with trypsin, and the resultant tryptic peptides were resolved by microbore reverse-phase HPLC (See Fig. 1B).

## EXAMPLE 2

In this example, peptide sequencing and cDNA cloning of the LAT protein is described. Peptides from five HPLC fractions were either microsequenced or subjected to mass spectrometry. The peptide from fraction 40 had a molecular weight of 1721.9 daltons (See Figure 1C). The residues at position 1, 2, 3, 11 and 15 could not be assigned unambiguously. The available sequence was then used to search the Genbank data base with the BLAST algorithm, and an EST clone from human fetal heart was found to encode the peptide fragment. Review of the Edman sequence data at the ambiguous positions was consistent with the EST sequence. The predicted molecular weight of the putative tryptic fragment predicted from the EST sequence was 1641.8 daltons. The 80.1 daltons difference suggested that a phosphate moiety of this molecular weight could be present at the Tyr residue. Two peptides from peaks 82 and 31 were shown by sequence analysis to be derived from the known tyrosine kinase substrate SLP-76. It was not clear whether these fragments of this 76kD substrate were cleaved physiologically or during protein purification. To avoid sequencing other peptides derived from SLP-76, the masses of subsequent peptides were determined. With this strategy, it was concluded that the peptide in fraction 55 was likely to originate from SLP-76, but the masses of two peptides in fraction 48 could not be of SLP-76 origin. The sequence of the larger of these two peptides was not found in the EST clone or Genbank. Interestingly, fraction 48 contained a small amount of a second peptide with a molecular weight of 1641.6, 80.3 daltons less than the mass of the peptide isolated and sequenced from peak 40. A partial amino acid sequence revealed that this peptide was the non-phosphorylated form of the peak 40 peptide.

A cDNA library from YT cells was probed with the *Eco* RI/ *Pst* I fragment of the EST clone. Thirty positive clones were isolated, of which the longest was 1.6 kb.

Determination of the nucleotide sequences of these clones predicted a protein that contains the two tryptic peptide sequences found in the isolated peptides (underlined in Fig.1D). Through comparison of the EST and cDNA sequences, it was found that the first 380 nucleotides in the EST clone were not present in the cDNA. This unknown sequence might be derived from

intronic sequence. Probes from the human cDNA were used to screen a murine adult thymus library. The murine amino acid sequence obtained was 66% identical to the human sequence (Fig.7D).

The coding sequence of human p36-38 began at nucleotide 58. The first methionine, not found within a consensus Kozak sequence, was followed by an open reading frame predicting a 233 amino acid polypeptide with a calculated molecular mass of 24.985 Daltons. One cDNA clone had an 87 nucleotide insertion at position 396, encoding an additional 29 amino acids. Of the 233 amino acids found in the majority of clones, there were a total of 39 negatively charged residues (16 Asp and 23 Glu), but only 11 positively charged residues (2 Lys and 9 Arg). The high relative negative charge could have resulted in retarded migration on SDS-PAGE leading to an apparent molecular weight of 36-38kD, in excess of the predicted mass. The deduced amino acid sequence contained no domain homologous to known tyrosine, serine/threonine, or lipid kinases, nor did it contain SH2, SH3, or PTB domains. A single region of hydrophobic amino acids extended from residues 5 to 27, and may form an  $\alpha$ -helical transmembrane domain. Two of the four residues N-terminal to this region were negatively charged in both human (two Glu) and murine (one Glu and one Asp). Positively charged residues (one Arg in the human and two Arg in the murine sequence), were found immediately after this hydrophobic domain. The relative charge difference across this putative transmembrane domain may determine the orientation of the molecule in the membrane, resulting in an extremely short extracellular amino-terminus and a long cytosolic tail.

Consistent with the prominent tyrosine phosphorylation of the protein in activated cells, the predicted cytosolic domain of human p36-38 contained ten tyrosines, of which nine were conserved between the human and murine proteins. Based on studies of Songyang and colleagues (Songyang *et al.*, Specific motifs recognized by the SH2 domains of Csk, 3BP2, fps/fes, GRB-2, HCP, SHC, Syk and Vav. *Mol. Cell. Biol.* 14, 2777-2785, 1994), after phosphorylation there were five potential binding sites for Grb2 SH2 domains (YxN) at Tyrosines 110, 127, 171, 191 and 226. Tyr171 and Tyr191 have an identical Grb2-binding motif (YVNV). A binding motif for the SH2 domain of the p85 subunit of PI3 kinase was not observed. A sequence related to the consensus binding motif for N-terminal and C-terminal PLC- $\gamma$ 1 SH2 domains (YLVV, Tyr132) were found to be present. Since the p36-38 protein most likely functioned as a docking protein capable of recruiting signaling

molecules, and in view of the observed features of its sequence. the name LAT for linker for activation of T cells was given.

To confirm that the cDNA clone encodes the full length LAT protein, and to resolve the discrepancy between the predicted and apparent molecular weight, the human LAT cDNA was subjected to *in vitro* transcription and translation. The reaction product contained a 38 kDa band that was not present when control DNA was used, confirming that the complete LAT coding sequence was present (not shown). The LAT cDNA was modified to include an epitopic tag (myc) at the C-terminus, cloned into the pcDNA3 expression vector and transfected into 293T cells (See Figure 2A). Immunoprecipitation and blotting demonstrated a 40kD protein in the transfected, but not non-transfected cells. These results also indicated that the LAT cDNA clone contained the entire coding region of the protein.

To demonstrate the LAT protein in T cells, rabbit polyclonal antibodies were raised against the cytosolic portion of LAT fused to GST. The resulting antiserum was used to detect LAT in lysates of activated Jurkat cells ( See Figure 2B). The LAT antiserum precipitated and detected a doublet of 36 and 38kD, which comigrated with bands detected with antiphosphotyrosine antibodies. Repetitive immunoprecipitation with this antiserum depleted most of p36-38 proteins from stimulated Jurkat lysate (data not shown). Phosphorylation of LAT partially interfered with the detection of this protein as demonstrated by enhanced blotting following dephosphorylation of the immunoprecipitate with calf intestinal phosphatase (CIP). This antiserum enabled the determination that LAT was rapidly tyrosine phosphorylated upon stimulation with OKT3 (See Figure 2C). Maximal phosphorylation of LAT was seen after stimulation for 15 seconds, and after 2 min LAT was rapidly dephosphorylated. These phosphorylation kinetics were also seen in whole cell lysates (not shown). The results in this example demonstrate that the LAT cDNA encodes a protein that migrates on SDS-PAGE with an apparent molecular weight of 36-38kD, and which is tyrosine phosphorylated, following TCR activation.

### EXAMPLE 3

In this example, tissue and cellular distribution of the LAT protein is described. Northern blot analysis was performed with poly (A)<sup>+</sup> RNAs isolated from different adult human tissues. Two transcripts (2.0 and 1.6 Kb) were seen predominantly in thymus, peripheral blood, and at a low level of expression in spleen. There were no transcripts of LAT found in other tissues (See Fig. 3A and data not shown). LAT expression was assayed

The cellular localization of LAT was analyzed by immunofluorescence in COS cells, which were transiently transfected with LAT cDNA, fixed, permeabilized and incubated sequentially with anti-LAT and TRITC-conjugated goat anti-rabbit antibodies. Visualization with confocal microscopy showed that LAT was localized to the plasma membrane, and a juxtanuclear intracellular compartment (data not shown). No staining was detected in untransfected COS cells. Plasma membrane localization of endogenous LAT was also specifically detected in Jurkat cells. When cells were simultaneously stained with anti-LAT and anti-Lck antibodies, LAT and Lck were seen to co-localize at the plasma membrane and in an intracellular compartment.

## 15

20

25

34

Syk, and that tyrosines in LAT involved in binding to the Grb2 SH2 domain are phosphorylated by these PTKs.

### EXAMPLE 5

5 In this example, association of LAT with Grb2, Grap, PI3K and PLC- $\gamma$ 1 is shown in 293T cells. To investigate the association of LAT with critical signaling molecules. FLAG-tagged LAT in the pcDNA3 vector was cotransfected in 293T cells with plasmids expressing HA-tagged Grb2, Myc-tagged Grap, HA-tagged p85 subunit of PI3K or PLC- $\gamma$ 1, in the presence or absence of both ZAP-70 and Lck. LAT was immunoprecipitated with  
10 anti-FLAG antibody, and associated proteins were detected by blotting with specific antibodies. As shown in Figure 4B, LAT tyrosine phosphorylation and association with endogenous Grb2 was induced by cotransfection with ZAP-70 and Lck as in Fig. 4A. When LAT and the two PTKs were coexpressed with HA-tagged Grb2, both endogenous Grb2 and HA-Grb2 bound to phosphorylated LAT (HA-Grb2 migrates more slowly than endogenous  
15 Grb2 on SDS-PAGE). Myc-tagged Grap, the Grb2-like protein, when coexpressed with LAT in the presence of the two PTKs, also associated with phosphorylated LAT. However, in this immunoprecipitation no endogenous Grb2 was bound to phosphorylated LAT, suggesting that Grap competes for Grb2 binding sites. HA-tagged p85 or PLC- $\gamma$ 1 also associated with phosphorylated LAT. Some endogenous Grb2 bound to LAT, but less than when LAT was  
20 cotransfected only with ZAP and Lck. Surprisingly, some association of LAT with PLC- $\gamma$ 1 was found, without co-transfection of the PTKs, suggesting either binding to undetectable tyrosine phosphorylation sites or binding via an SH2-independent mechanism.

The above experiments were designed to demonstrate LAT associations by isolation of LAT followed by detection of associated molecules. The reciprocal approach was also  
25 performed to confirm these associations. 293T cells were co-transfected with FLAG-tagged LAT, ZAP-70 and Lck, and either HA-tagged Grb2 or myc-tagged Grap (Fig.4C). As above, immunoprecipitation of LAT allowed detection of that fraction of Grb2 or Grap associated with LAT. In addition, immunoprecipitation of either Grb2-HA or Grap-myc allowed detection of the fraction of LAT bound to either adaptor protein. Similar experiments were  
30 performed in cells co-transfected with HA-p85 or PLC- $\gamma$ 1. Immunoprecipitation of these proteins revealed associated LAT (not shown). Additional experiments were performed to determine if the PTKs, ZAP-70 or Lck, could bind to LAT. No associations between LAT and either PTK were detected (not shown).

## EXAMPLE 6

In this example, association of LAT with Grb2, PLC- $\gamma$ 1 and PI3K is shown in Jurkat T cells. The ability of LAT to associate with signaling proteins in T cells was next evaluated. The interaction of Grb2 with p36-38 is thought to be mediated by the Grb2 SH2 domain. To test this, lysates prepared from resting and activated Jurkat cells stimulated with C305 were incubated with the immobilized fusion proteins, GST-Grb2, GST-Grb2 with an SH2 domain mutation (R86K), and GST-Grb2 with mutations at both SH3 domains (P49L/G203R). Only GST-Grb2 and GST-Grb2 with SH3 mutations bound a 36-38 kDa tyrosine phosphorylated protein from C305 stimulated Jurkat lysate (Figure 5A). Mutation at the SH2 domain of Grb2 abolished the interaction between this 36-38 kDa protein and GST-Grb2. Subsequently the same membrane was probed with anti-LAT antibodies which confirmed that this p36-38 is LAT. LAT was not detected in material subjected to purification with GST-Grb2 from unstimulated cells or with GST-Grb2 containing the SH2 mutation.

To demonstrate the interactions in T cells directly, lysates from unstimulated and stimulated Jurkat T cells were subjected to immunoprecipitation with anti-Grb2 and anti-PLC- $\gamma$ 1 antibodies followed by blotting with anti-LAT. As shown in Fig. 5B, equal amounts of Grb2 or PLC- $\gamma$ 1 were immunoprecipitated from unstimulated or stimulated lysates. When the membrane was blotted with anti-phosphotyrosine antibody, a 36-38 kDa tyrosine phosphorylated protein was detected in association with Grb2 and PLC- $\gamma$ 1 only in activated T cells. This protein was identified as LAT by immunoblotting with anti-LAT antibody. CIP treatment of the immunoprecipitate enhanced LAT detection. p85 in anti-Grb2, but not in anti-PLC- $\gamma$ 1 immunoprecipitates in lysates of activated Jurkat cells, was also detected. There was also some low level of association between Grb2 and PLC- $\gamma$ 1 probably through LAT (data not shown).

TCR-mediated activation results in tyrosine phosphorylation of many substrates (Figs. 1A and 4C). To identify additional proteins, capable of binding LAT following T cell activation, anti-LAT antisera was used to immunoprecipitate LAT from lysates of resting and activated Jurkat cells. As shown in Fig. 5C, several tyrosine phosphorylated proteins with molecular weights of about 70, 76, 100, 120, 135 kD were specifically co-precipitated with anti-LAT only from lysates of stimulated Jurkat cells. By using a panel of antibodies to immunoblot known tyrosine kinase substrates, some of these proteins were identified as PLC- $\gamma$ 1 (135kD), Cbl (120 kD), Vav (100 kD) (not shown). The 76kD protein was most likely SLP-76. The associations of SLP-76 and Cbl with LAT were probably mediated

indirectly through Grb2, because both bind the Grb2 SH3 domain. Vav was likely detected because of its association with SLP-76. The 70 kDa protein was not identified, and was not Sam68 or ZAP-70. Anti-LAT immunoprecipitates from stimulated Jurkat also contained the p85 subunit of PI3K and Grb2 (not shown). The results in this example, demonstrate that LAT is present in signaling complexes containing multiple critical molecules.

### EXAMPLE 7

In this example, the function of LAT in the TCR signal transduction pathway is shown. The hypothesis from the above studies is that the association of tyrosine phosphorylated LAT with the various signaling molecules is required for the TCR signal transduction process. To directly address the role of LAT in signaling through the TCR, wild type LAT or a mutant form of LAT was overexpressed and the effect of this overexpression examined on TCR-mediated activation. The mutant form of LAT employed in these experiments contains Phe for Tyr substitutions at both positions Y171 and Y191 (Y171/191F). These two tyrosine residues were chosen for mutagenesis because they are within identical YVNV motifs, a motif which has previously been shown to mediate the binding of Shc to Grb2 (See Songyang *et al.*, Specific motifs recognized by the SH2 domains of Csk, 3BP2, fps/fes, GRB-2, HCP, SHC, Syk and Vav. *Mol. Cell. Biol.* 14, 2777-2785, 1994). Moreover, phosphorylation at Tyr 191 was detected during microsequencing and mass spectrometry analysis of peptides from p36-38 (see above).

To investigate the effect of mutant LAT on protein-protein interactions, stable cell lines overexpressing myc-tagged wild type or Y171/191F LAT were established. Myc-tagged LAT was immunoprecipitated with anti-myc antibody from unstimulated or C305 stimulated transfectants. As shown in Fig. 6A, blotting with anti-LAT antibodies demonstrated that the amount of LAT immunoprecipitated with anti-myc was comparable in the two cell lines. The myc-tagged LAT appears as a doublet as does endogenous LAT. WT and Y171/191F LAT were both tyrosine phosphorylated upon stimulation, though the level of Y171/191F tyrosine phosphorylation was less than that of WT LAT, suggesting that tyrosine residues besides Y171 and Y191 are phosphorylated. Though p85, Grb2 and PLC- $\gamma$ 1 were observed to bind the WT form of LAT, mutations at Y171 and Y191 abolished the binding of Grb2 and p85, and greatly reduced the binding of PLC- $\gamma$ 1 upon activation. In addition, mutations at Y171 and Y191 prevented the association of SLP-76, Vav and Cbl, but the 70 kDa tyrosine-phosphorylated protein remained associated with LAT. Additionally, the Ras

activator protein SOS was observed to co-precipitate with LAT, likely due to its interaction with Grb2. The association with SOS was also disrupted by the double Y to F mutations.

The functional effect of mutant LAT overexpression was assayed by determining the effect on transcriptional activation of AP-1 and NF-AT, both involved in TCR-mediated transcriptional events. Jurkat TAg cells, were transiently transfected with wild-type LAT, mutant LAT(Y171/191F), or vector only, together with a reporter construct for secreted alkaline phosphatase (SEAP) driven by the AP-1 or NF-AT response elements. Fig 6B demonstrates that overexpression of WT LAT did not result in any significant effect on AP-1 transcriptional activity and only slightly increased NF-AT transcriptional activity compared with transfection of the control vector (see Fig 6). However, overexpression of the mutant form of LAT (Y171/191F) blocked TCR-mediated AP-1 and NF-AT transcriptional activity. The results in this example indicates that LAT plays an important role in activation of transcription mediated by AP-1 and NF-AT following TCR stimulation.

From the above, it should be clear that the present invention provides methods and compositions useful for identifying signaling pathway agonists and antagonists. The methods and compositions are, in this manner, useful for identifying compounds that may be used *in vitro* and *in vivo* in the context of T cell, NK cell and mast cell responses.

### EXAMPLE 8

In this example the discovery of LAT palmitoylation sites and the use of LAT and LAT with modified palmitoylation sites in the TCR signal pathway is shown. Human and murine LAT contain two conserved cysteines, C26 and C29, near the punitive transmembrane domain. These features suggest that LAT could be a member of a class of transmembrane proteins modified by palmitate at juxamembrane cysteine residues. To examine whether LAT is palmitoylated, Jurkat cells were metabolically labeled with [<sup>3</sup>H]palmitate. LAT was immunoprecipitated with anti-LAT antiserum, and immunoprecipitates were analyzed by SDS-PAGE and flouorography. [<sup>3</sup>H]palmitate was found to be incorporated into LAT. When an identical gel was treated with 1M hydroxylamine, the [<sup>3</sup>H] signal was almost completely lost, indicating that the addition of palmitate to LAT was through S-acylation.

Many palmitoylated proteins, such as the Src family kinases Lck and Fyn, are targeted into glycolipid-enriched microdomains (GEMs) of the plasma membrane (Alland *et al.* "Dual myristylation and palmitoylation of Src family member p59<sup>fyn</sup> affects subcellular localization" *J. Biol. Chem.* 269:16701-16705, 1994; Shenoy-Scaria *et al.* "Cysteine of Src family protein



tyrosine kinases determines palmitoylation and localization in caveolae" *J. Cell Biol.* 126:353-363, 1994). GEMs are operationally characterized by resistance to Triton extraction at 4°C and are found in low-density gradients of a sucrose gradient (Simons and Ikonen "Functional rafts in cell membranes" *Nature* 387:569-572, 1997). Since LAT is palmitoylated, we next examined whether LAT localizes to GEMs. Jurkat cells were solubilized with 1% Triton, and lysates were subject to overnight ultracentrifugation over a sucrose gradient (5%/30%/40% step gradient). Among a total of 12 fractions collected, fraction 3, at the interface between 5% sucrose and 30% sucrose, contained the low-density membrane fraction including the GEMs (Brown and Rose "Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface" *Cell* 68:533-544, 1992). Fractions 8-12 from the 40% sucrose section were the Triton-soluble fractions. LAT was detected by immunoblotting and was found in both GEM and Triton-soluble fractions. LAT in the GEM fractions (e.g. fraction 3) was predominately the p36 form. Two prominent forms of LAT, p36 and p38, were detected in Triton-soluble fractions (e.g. fraction 10). LAT was more abundant in fraction 3 than in fraction 10 as determined by immunoblot. When a sample of fraction 3 was diluted 16-fold, the amount of LAT still exceeded that found in the undiluted fraction 10. As controls for GEM purification, we immunoblotted the same membrane with anti-Lck and anti-ZAP70 antibodies. Lck was observed in both GEM and Triton-soluble fractions, but ZAP-70 was found only in Triton-soluble fractions, as expected.

To study the relationship between palmitoylation and GEM localization, Jurkat cells were labeled with [<sup>3</sup>H]palmitate for 3 hr, lysed in 1% Triton, and subject to ultracentrifugation on a sucrose gradient. LAT was then immunoprecipitated from GEM and Triton-soluble fractions, resolved by SDS-PAGE, and analyzed by both fluorography and anti-LAT immunoblotting. The anti-LAT immunoblot showed that an approximately equal amount of LAT was precipitated from these two fractions. However, LAT in the GEM fraction was clearly more palmitoylated, indicating that palmitoylated LAT predominately localizes to GEMs.

Since LAT is present in both GEM and Triton-soluble fractions, we next examined in which fraction tyrosine phosphorylated LAT can be found after stimulation of Jurkat cells. We also studied whether activation of T cells induces redistribution of other signaling molecules into GEMs. Crosslinking with the anti-CD3ε antibody, OKT3, leads to tyrosine phosphorylation of many cellular proteins and activation of T cells. Unstimulated or OKT3-stimulated Jurkat lysates were subjected to sucrose gradient centrifugation to separate GEM

fractions from Triton-soluble fractions. Each fraction was analyzed by SDS-PAGE followed by immunoblotting with anti-phosphotyrosine antibody and antibodies against each individual protein. As shown in Fig. 13A, LAT and Lck were basally tyrosine phosphorylated in both GEM and Triton-soluble fractions from unstimulated cells. A tyrosine phosphorylated 80 kD diffuse band was also observed in the GEM fractions. LAT phosphorylation results in interactions with Vav, a membrane associated guanine nucleotide releasing factor. While the majority of Vav localized to the Triton-soluble fractions, some tyrosine phosphorylated Vav was detected in the GEM fractions (Fig.13A) and fractions 5-7, as shown by anti-Vav blot in Fig.13C.

Upon stimulation with OKT3, most tyrosine phosphorylated LAT appeared in the GEM fractions, although some was in the Triton-soluble fractions (Fig. 13B). There was no obvious change in Lck tyrosine phosphorylation, but the 60 kD form of Lck, well known to accompany TCR activation (Marth *et al.* "Lymphocyte activation provokes modification of a lymphocyte-specific protein tyrosine kinase (p56lck)" *J. Immunol.* 142:2430-2437, 1989), was increased in Triton-soluble fractions upon stimulation, as shown by the anti-Lck blot (Fig. 13C). Other prominent tyrosine phosphorylated proteins present in the GEM fractions upon stimulation were PLC- $\gamma$ 1 and Vav (Fig. 13B). Blotting with antibodies to particular signaling molecules supported the conclusion that certain molecules re-distribute to GEMs upon TCR activation (Fig. 13C). The amount of PLC- $\gamma$ 1 increased in the GEM fractions and fractions 5-7 upon stimulation. Like PLC- $\gamma$ 1, the amount of Vav in those fractions also increased after stimulation. The distribution of the p85 subunit of PI-3 kinase was similar to Vav. There was also some increase of Grb2 in the GEM fractions after stimulation (Fig. 13C). Cbl and ZAP-70 were present in Triton-soluble fractions, but they did not re-distribute into GEMs after stimulation (Fig.13C). We used the detergent octyl-glucoside to solubilize GEMs and Triton-soluble fractions in order to assess LAT-associated proteins in the two pools. This detergent is capable of solubilizing Triton-insoluble GEMs (Brown and Rose "Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface" *Cell* 68:533-544, 1992). Following extraction with this detergent, we observed by co-immunoprecipitation that more PLC- $\gamma$ 1 and Vav associated with LAT in GEMs than LAT in Triton-soluble fractions, suggesting that LAT in GEMs interacts with these signaling molecules following activation.

The present invention contemplates assays for the screening of compounds that are agonistic or antagonistic for LAT activation or that are agonistic or antagonistic for LAT

binding of associated molecules. For example, but not by way of limitation, it is contemplated that compounds will be screened using LAT, or portions thereof, and the methods disclosed herein (*e.g.* determining the extent of LAT incorporation into GEMs). In such an assay compounds that activate LAT would result in a greater amount of LAT being associated with GEMs, as compared to controls. Likewise, compounds that interfered with LAT activation would result in lesser amounts of LAT being associated with GEMs, as compared to controls. Additionally, but not by way of limitation, it is contemplated that compounds that modulate LAT binding to associated molecules (for example, but not limited to, Vav, PLC- $\gamma$ 1, Grb2, Sos, cb1, SLP-76, SLAP, and PI-3 kinase) will be screened using LAT, or portions thereof, and the methods disclosed herein. Modulation of LAT binding of any of these molecules can be determined by methods known to those practiced in the art (*e.g.* immunoblot, measurement of binding coefficients, measurement of downstream activation events).

To test whether two cysteines, C26 and C29, in human LAT are the sites for addition of the palmitate moiety, we made myc-tagged LAT mutants with single cysteine to alanine mutations (C26A, C29A) and a double mutant (C26/29A). We also made a mutant LAT ( $\Delta$ tm) with a deletion of the N-terminal region (residues 1-22) including the transmembrane domain, to investigate the role of the transmembrane domain in LAT palmitoylation and function. T-antigen transformed human kidney fibroblasts (293T cells) were transiently transfected with these constructs and metabolically labeled with [ $^3$ H]-palmitate. LAT was then immunoprecipitated with anti-myc antibody from lysates of these transfected cells. As seen in Fig.14 (top panel), two forms of LAT labeled with [ $^3$ H]-palmitate were detected in immunoprecipitates of wild type (wt) LAT. For the C26A mutant, only the p38 form was labeled with [ $^3$ H]-palmitate and the amount of  $^3$ H label in C26A was less than in wt. Two forms of LAT labeled with [ $^3$ H]-palmitate were also observed with the C29A mutant, but compared with wt, the amount of  $^3$ H incorporated into the p38 form was reduced and  $^3$ H label in the p36 form was barely detectable. A similar trace amount of  $^3$ H was present in the LAT double cysteine mutant (C26/29A) and in the transmembrane domain deletion mutant ( $\Delta$ tm).

The same samples from the above labeling experiment were loaded on another gel and analyzed by blotting with anti-myc antibody. As shown in Fig.14 (lower panel), similar amounts of proteins were expressed in each lane. Note that the only the p314 form of LAT was detected by blotting indicating in transient transfections the p36 form was rare. These results indicate that C26 and C29 are the major sites for LAT palmitoylation. The trace

amount of  $^3\text{H}$  in C26/29A and  $\Delta\text{tm}$  mutant LAT is most likely due to metabolic conversion of  $^3\text{H}$ -palmitate into  $^3\text{H}$ -amino acids, although we could not totally exclude the possibility that LAT might be palmitoylated at a low level at other site(s).

Because mutations of two cysteines, C26 and C29, affected the palmitoylation of LAT, we next tested whether these mutations also altered membrane localization and the partition of LAT into GEMs. We examined the membrane localization of wt and mutant LAT by immunofluorescence staining of transiently transfected 293T cells. There was no significant difference in localization between wt LAT and any of the cysteine mutants. However, deletion of the transmembrane domain of LAT resulted into a predominantly cytosolic localization (Fig. 15A). We also transiently transfected Jurkat/TAg cells (SV40 large T antigen transformed Jurkat cells, Clipstone and Crabtree "Identification of calcineurin as a key signaling enzyme in T-lymphocyte activation" *Nature* 357:695-697, 1992) with LAT constructs and analyzed the subcellular distribution of LAT by fractionation of transfected cells into cytosolic and particulate fractions. Myc-tagged wt LAT, C26A, C29A, and C26/29A mutants predominantly localized to the particulate fraction, while the LAT  $\Delta\text{tm}$  was detected mainly in the cytosolic fraction (Fig. 15B).

Although C26A, C29A, and C26/29A LAT mutants all localized to the plasma membrane, it was possible that these mutations might affect the targeting of LAT into GEMs. Transiently transfected Jurkat cells were lysed in 1% Triton, and GEM fractions were purified using a sucrose gradient. Samples from fractions 3 and 10, as representatives of the GEM and Triton-soluble fractions, respectively, were analyzed on SDS-PAGE and immunoblotted with anti-myc antibody. The wt LAT and all of the LAT mutants were expressed equally in Triton-soluble fractions (Fig. 15C). Two forms of wt LAT and C29A were observed in these fractions, though the p36 form of C29A was barely detectable. Strikingly, only the wt LAT and a small amount of C29A were present in the GEM fractions. The plasma membrane localization of C26A and C26/29A and the palmitoylation of the C26A mutant were not sufficient for GEM targeting. The cytosolic LAT  $\Delta\text{tm}$  mutant was not seen in the GEM fraction, as expected. These results indicate that, while the transmembrane domain alone suffices for LAT plasma membrane localization, palmitoylation at C26 is essential for GEM targeting. Palmitoylation only at C29 is not sufficient for GEM targeting, but it does increase the efficiency of C26-mediated targeting.

We next addressed whether palmitoylation and targeting into GEMs were necessary for the phosphorylation of LAT by tyrosine kinases. Jurkat cells were transfected with

mutant LAT constructs and stimulated with OKT3 at 36 hours after transfection. The p36 form of wt myc-tagged LAT was rapidly tyrosine phosphorylated upon OKT3 stimulation (Fig.10). The p36 form of the C29A mutant was also tyrosine phosphorylated, though less than wt. Only very weak tyrosine phosphorylation of the p314 form of C26A or C26/29A was observed. Surprisingly, LAT  $\Delta$ tm was also tyrosine phosphorylated to some extent. This could be explained by the presence of activated ZAP-70 in the cytosol following stimulation. These activated ZAP-70 molecules might phosphorylate LAT  $\Delta$ tm in the cytosol. Endogenous LAT, however, must be in GEMs to become optimally tyrosine phosphorylated during activation. Blotting the same membrane with anti-myc antibody showed that the amount of LAT was nearly equal in all lanes. This protein blot also revealed the presence of a band at 140 kD in the wt and C29A lanes. This band could be detected with anti-LAT antibody. This 140 kD band could be a dimer of LAT, resistant to treatment with SDS and DTT. In conclusion, our results clearly show that LAT palmitoylation at C26 and C29 not only is required for LAT palmitoylation, but also is necessary for efficient tyrosine phosphorylation by tyrosine kinases during T cell activation.

The present invention contemplates the use of these LAT mutations (C26A, C29A, C26/29A and  $\Delta$ tm), or parts thereof, in screening assays for compounds that are agonistic or antagonistic to LAT activation. Additionally, the present invention contemplates the use of these LAT mutations in screening assays for compounds that will allow for the activation of T cells in the presence of non-functional (*e.g.* non-palmitoylated) LAT.

### EXAMPLE 9

In this example the staining of both cryostat and paraffin embedded normal tissues and cells was performed. Positive staining for LAT was crisp and mainly localized on the cell membrane and the sub-plasmalemmal area. Variable number of cells also showed diffuse cytoplasmic reactivity and, on occasion, dot-like positivity in the Golgi region. On paraffin sections, optimal staining was observed in tissues fixed in both buffered formalin or B5, whereas occasionally some background staining was observed for tissues fixed in Bouin or Hollande. Decalcification of bone marrow specimens in EDTA did not affect LAT immunoreactivity.

In the thymus, expression of LAT was identifiable throughout all stages of thymocyte differentiation, including the large cortical blasts. In embryos of 10-12 weeks of gestation, thymuses already showing a lobular architecture contained lymphoid cells that expressed LAT

and CD3, but were negative for TdT indicating that LAT expression precedes TCR rearrangement.

In peripheral lymphoid tissues, LAT-positive lymphocytes were located in the known T cell areas in lymph nodes and spleen, and the immunoreactivity paralleled that obtained with anti-CD3 in both frozen and paraffin sections. In the small intestine, intraepithelial T cells were also positive for LAT. In bone marrow, LAT was expressed by the sparse T lymphocytes present in interstitial spaces, and also by platelets and megakaryocytes, that exhibited a strong reactivity in the cytoplasm; all other hematopoietic cells were negative. Reactivity for LAT was also noticed on tissue mast cells, in the form of delicate plasma membrane and granular cytoplasmic labeling. In all tissues, non-T cell components, including B cells, macrophages, plasmacytoid monocytes, epithelioid histiocytes and dendritic cells, were completely negative for LAT. Similarly, LAT was not expressed on endothelial cells and epithelia, including thymic epithelium and Hassal bodies.

On peripheral blood mononuclear cells (PBMC), LAT was expressed, in addition to CD3+ T cells, on resting CD16+ and CD56+ NK cells. LAT staining was also found on T and NK cells in culture with rIL-2; on CD3-CD16+CD56+ NK cells purified from PBMC, LAT was obviously expressed, although the intensity of reactivity was much weaker than that recognized on purified CD3+ T cells.

#### EXAMPLE 10

In this example the staining of LAT in neoplastic tissues was performed. Neoplastic cells in all cases of Hodgkin's (15 cases) and B cell non-Hodgkin's lymphomas (100 cases) (see Table 1) were negative for LAT, but contained variable amounts of LAT-reactive non neoplastic T cells. In T-cell lymphomas (excluding anaplastic large cell lymphomas (ALCL)), LAT stained 14/14 lymphoblastic lymphomas, 42/43 peripheral T cell lymphomas, and 11/12 NK/T cell lymphomas. The intensity of reactivity and the percentage of labeled cells varied among different types of T cell lymphomas. In lymphoblastic lymphomas all neoplastic cells were strongly labeled by anti-LAT; in contrast, in peripheral T cell and NK/T cell lymphomas the reactivity was intense in 214/53 (52.14%) cases, and it was recognizable in the vast majority of cells in 50/53 (94.3%). LAT expression showed no substantial differences in peripheral T cell lymphomas with relationship to primary location, cell morphology, and immunophenotype, with the exception of NK/T cell lymphomas, which were more frequently characterized by weaker and a lower number of labeled cells.

**Table 1.**

**Details of negative LAT staining Hodgkin's lymphomas and malignant B cell lymphomas analyzed.**

	<b>N° of cases</b>
<b>Hodgkin's Lymphomas</b>	
Lymphocyte predominance	2
Nodular sclerosis	10
Mixed cellularity	2
<b>Total</b>	<b>15</b>
<b>B-cell lymphomas</b>	
I. Precursor B-lymphoblastic leukemia/lymphoma	8
II. Peripheral B-cell neoplasms	
B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma	14*
Lymphoplasmacytoid lymphoma	3
Mantle cell lymphoma	11**
Follicle center cell lymphoma, follicular	16
Marginal zone B-cell lymphoma	
nodal	1
extranodal	4
Hairy cell leukemia	6
Myeloma	8
Diffuse large B-cell lymphoma	25***
Burkitt's lymphoma	4
<b>Total</b>	<b>100</b>

\*: one with immunoblastic transformation; \*\* one "blastoid" variant; \*\*\*: included four T-cell rich B large cell lymphomas and three CD30+/ALK1- anaplastic large cell lymphomas.

The ALCL included 39 extra-cutaneous and 5 purely cutaneous cases: on the basis of the expression of at least one T cell marker (CD3, CD5, CD14, CD43, CD45R0, TCR- $\beta$ ) and no B cell markers (CD20 or CD79a), 25 cases were classified as T cell and 19 cases as null-cell phenotype. 14 cases (31.14%) reacted with LAT; all were represented by extra-cutaneous lymphomas, ten of them were of T cell phenotype, while four were null-cell. LAT positive ALCL showed intense reactivity in 5/14 (35.7%) cases, and it was recognizable in the vast majority of cells in 10/14 (71.4%).

LAT and CD3 expression was compared in T cell and NK/T cell lymphomas (excluding ALCL). In T-lymphoblastic lymphoma, LAT positivity was more intense and diffuse than CD3. Among peripheral T cell lymphomas, only two LAT-negative cases were observed. One was a large cell lymphoma presenting an aberrant phenotype with loss of CD3, CD5, CD43 and TCR- $\beta$ , the other LAT negative case was a NK/T which strongly expressed cytoplasmic CD3. Three cases of LAT positive cutaneous T-cell lymphomas did not express CD3 (two also lacked CD5), but were positive for CD2, CD4 and TCR- $\beta$ .

A comparison was made regarding LAT, CD3 and ALK1 expression in ALCL. The distribution of LAT, CD3 and ALK1 in ALCL is reported in Table 2. Seven ALCL cases were LAT+CD3+, 7 were LAT+ CD3-, 9 were LAT- CD3+, and 21 were LAT-CD3-. Among ALCL T cell type (25 cases), CD3 labeled a higher number of cases (19/25; 76%) than LAT (10/25; 40%). Interestingly, LAT and CD3 expression in ALCL T cell type was discordant in a significantly higher number of cases than in other peripheral T cell lymphomas (12/25 versus 4/55, respectively; Fisher's exact test:  $p < 0.001$ ). Immunostaining for ALK1 was available in 42 cases and showed nuclear  $\pm$  cytoplasmic positivity in 10. Although the majority of ALK1+ cases were LAT+ (6/10), in comparison with ALK1- cases (14/32), this difference was not statistically significant (Fisher's exact test:  $p = 0.059$ ).

Expression of LAT was also examined in Hematopoietic, non-lymphoid neoplasms. In chronic myeloproliferative disorders and myelodysplastic syndromes, cytoplasmic LAT expression was consistently identified in normal and atypical megakaryocytes, including the micro-megakaryocytes typically found in chronic myeloid leukemia and myelodysplasia. Among 214 cases of acute non-lymphoid leukemia, the only positive cases were represented by two megakaryocytic leukemias, where the blast cells were labeled in their cytoplasm; an additional case of chronic myeloid leukemia in blastic transformation contained two cell populations, one formed by myeloid blasts (myeloperoxidase and CD34 positive), another by megakaryocytic blasts, expressing LAT, Factor VIII-related antigen and CD34. In all



Table 2. Details of LAT, CD3 and ALK1 staining in anaplastic large cell lymphomas (ALCL).

	T-cell type				Null-cell type			
	LAT+ CD3+	LAT+ CD3-	LAT- CD3-	LAT- CD3+	LAT+ CD3+	LAT+ CD3-	LAT- CD3-	LAT- CD3+
ALK1 positive	3	3	1		3			
ALK1 nega- tive	4		4	8	4      12			
ALK1 n.a.*			1	1				
Total	7	3	6	9	0	4	15	0

\*: ALK1 not assessable because only B5-fixed specimens available and poor reactivity of anti-ALK1 on this fixative.

pathological conditions. reactivity of LAT on megakaryocytes was strong and particularly helpful in the identification of abnormal forms; moreover, it was frequently stronger than that of other megakaryocytic markers on paraffin sections, such as Factor-VIII related antigen and CD61. 4/5 cases of mastocytosis showed LAT positivity, including 3 of 3 cases of cutaneous mastocytosis, and 1 of 2 cases of systemic mastocytosis, as with normal mast cells. LAT expression in neoplastic mast cells was finely granular in the cytoplasm and delicate on the cell membrane, a pattern of staining that clearly differed from that on T cells.

## CLAIMS

We claim:

1. An isolated polypeptide comprising at least a portion of the amino acid  
5 sequence of SEQ ID NO:4.

2. The isolated polypeptide of Claim 1, wherein said portion comprises a region  
comprising at least one tyrosine.

10 3. The isolated polypeptide of Claim 1, wherein said portion comprises a region  
defined by amino acids 28 to 233.

4. A purified antibody which binds specifically to a polypeptide comprising at  
least a portion of the amino acid sequence of SEQ ID NO:4.

15 5. The purified antibody of Claim 4, wherein said antibody is a polyclonal  
antibody.

6. The purified antibody of Claim 4, wherein said antibody is a monoclonal  
20 antibody.

7. An isolated polynucleotide encoding the polypeptide comprising the sequence  
of SEQ ID NO:4.

25 8. The isolated polynucleotide of Claim 7, wherein said polynucleotide comprises  
the sequence of SEQ ID NO: 1

9. The isolated polynucleotide of Claim 7, wherein said polynucleotide is  
contained on a recombinant expression vector.

30 10. The polynucleotide sequence of Claim 9, wherein said expression vector  
containing said polynucleotide sequence is contained within a host cell.

11. A polynucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO:1.

12. A method of screening a compound, said method comprising:

a) providing, in any order:

i) a peptide comprising at least a portion of the amino acid sequence set forth in SEQ ID NO:4, wherein said portion is capable of binding to a LAT binding ligand;

ii) a LAT binding ligand; and

iii) one or more compounds for screening;

b) mixing, in any order, said peptide, said LAT binding ligand and said one or more compound; and

c) measuring the extent of binding of said peptide to said LAT binding ligand.

13. The method of Claim 12, wherein said LAT binding ligand comprises a tyrosine kinase.

14. The method of Claim 13, wherein said kinase comprises ZAP-70 kinase.

15. The method of Claim 13, wherein said kinase comprises Syk kinase.

16. The method of Claim 12, wherein said peptide is part of a fusion protein.

17. A method for detecting the presence of a portion of the polypeptide having the amino acid sequence set forth in SEQ ID NO:4, said method comprising the steps of:

a) providing in any order:

i) an antibody capable of reacting with a portion of the polypeptide having the sequence set forth in SEQ ID NO:4; and

ii) a sample suspected of containing at least a portion of the polypeptide having the sequence set forth in SEQ ID NO: 4:

- b) combining said antibody and said sample under conditions such that a complex is formed between said antibody and said portion of said polynucleotide; and
- c) detecting said complex.

18. The method of Claim 17, wherein said antibody is a polyclonal antibody.

19. The method of Claim 17, wherein said antibody is a monoclonal antibody.

20. The method of Claim 17, wherein said sample comprises lymphocytes.

21. A method for detecting the presence of polynucleotide sequences encoding at least a portion of LAT gene in a sample, said method comprising the steps of:

a) providing in any order:

i) a polynucleotide comprising a sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO:1; and

ii) a sample suspected of containing nucleic acid comprising the sequence of SEQ ID NO:1;

- b) combining said polynucleotide and said sample under conditions such that a hybridization complex is formed between said polynucleotide and said sample nucleic acid; and
- c) detecting said hybridization complex.


22. The method of Claim 21, wherein said sample nucleic acid is RNA.

23. The method of Claim 21, wherein said sample nucleic acid is DNA.

24. The method of Claim 21, wherein said sample comprises lymphocytes.

25. The method of Claim 24, wherein said detected hybridization complex correlates with expression of the LAT gene in said lymphocytes.

C



Expt	Peak	Mass	Sequence	Source
1	40	1721.9	x x v n v s q e l h p x a a k	LAT
1	82	1840.0	s e v l g w d p d s l a d y f k	SLP-76
2	31	n.d.	s i p t r	SLP-76
3	55	1334.8	n.d.	SLP-76
3	48	1743.3	l p o s y d s t s s d s l y p r	LAT
3	48	1641.6	x y v n v	LAT

B




Figure 1

2/24






Figure 2

3/24

**A**






**B**




**Figure 3**





**Transfection with Lck, ZAP-70 & LAT-FLAG plus:**



### Figure 4

5/24




Figure 5

6/24




Figure 6

## Human LAT Nucleotide Sequence (1-1060)

1 gactctgccc ttgaggggcc taggggtgca gccagcctgc tccagctcc cctgcagatg  
61 gaggaggcca tcttggtccc ctgcgtgctg gggctcctgc tctgcccac cctggccatg  
121 ttgatggcac tgttgtgca ctgccacaga ctgccaggct cctacgacag cacatcctca  
181 gatagtttgt atccaagggg catccagttc aaacggcctc acacggttg cccctggcca  
241 cctgcctacc cacctgtcac ctctaccca cccctgagcc agccagacct gctcccatc  
301 ccaagatccc cgcagccct tgggggtcc caccggacgc catctcccg gcgggattct  
361 gatggtgcca acagtgtggc gagctacgag aacgaggaac cagcctgtga ggatgcagat  
421 gaggatgagg acgactatca caaccaggc tacctggtgg tcttctga cagcaccg  
481 gccactagca ctgctgccc atcagctct gcactcagca cccctggcat ccgagacagt  
541 gccttctcca tggagtccat tgatgattac gtgaacgttc cggagagcgg ggagagcgca  
601 gaagcgtctc tggatggcag ccgggagtat gtgaatgtgt cccaggaact gcacctgga  
661 gcggctaaga ctgagcctgc cgccctgagt tcccaggagg cagaggaagt ggaggaagag  
721 ggggctccag attacgagaa tctgcaggag ctgaactgag ggcctgtgga ggccgagtct  
781 gtcctggaac caggcttgcc tgggacggct gagctgggca gctggaagtg gctctgggt  
841 cctcacatgg cgtctgccc ttgtccagc ctgacaacag cctgagaaat cccccgtaa  
901 cttattatca ctttgggtt cggcctgtgt ccccgaacg ctctgcacct tctgacgag  
961 cctgagaatg acctgccctg gcccagccc tactctgtgt aatagaataa aggctgcgt  
1021 gtgtctgtgg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

Figure 7A

8/24

## Human LAT Nucleotide Sequence (1-1460), [alternative splice variant]

1 accccatctt catctggcct tgactctgcc cttgaggggc ctaggggtgc agccagcctg  
61 ctccgagctc cctgcagat ggaggaggcc atcctgggcc cctgcgtgct ggggctcctg  
121 ctgctgccc tctggccat gtgatggca ctgtgtgtgc actgccacag actgccaggc  
181 tctacgaca gcacatcctc agatagtttg tatccaaggg gcatccagtt caaacggcct  
241 cacacggttg cccctggcc acctgcctac ccacctgtca cctctaccc acccctgagc  
301 cagccagacc tgcctcccat cccaagatcc ccgcagcccc ttgggggtgc ccaccggacg  
361 ccattctccc ggcgggattc tgatggtgcc aacagtgtgg cgagctacga gaacgagggt  
421 gcgtctggga tccgaggtgc ccaggctggg tggggagtct ggggtccgtc ctggactagg  
481 ctgacctctg tctgttacc cccagaacca gcctgtgagg atgcagatga ggatgaggac  
541 gactatcaca acccaggcta cctggtggtg cttcctgaca gacccccggc cactagcact  
601 gctgccccat cagctcctgc actcagcacc cctggcatcc gagacagtgc ctctccatg  
661 gagtccattg atgattacgt gaacgttccg gagagcgggg agagcgcaga agcgtctctg  
721 gatggcagcc gggagtatgt gaatgtgtcc caggaaactgc atcctggagc ggctaagact  
781 gagcctgccg cctgagttc ccaggaggca gaggaagtgg aggaagaggg ggctccagat  
841 tacgagaatc tgcaggagct gaactgaggg cctgtggagg ccgagtctgt cctggaacca  
901 ggcttgccctg ggacggctga gctgggcagc tggaagtggc tctgggggtc tcacatggcg  
961 tctgcccctt gctccagcct gacaacagcc tgagaaatcc cccgtaact tattatcact  
1021 ttggggttcg gcctgtgtcc cccgaacgct ctgcacctc tgacgcagcc tgagaatgac  
1081 ctgccctggc cccagcccta ctctgtgtaa tagaataaag gcctgcgtgt gtctgtgtg  
1141 agcgtgcgtc tgtgtgtgcc tgtgtgcgag tctgagtcag agatttgag atgtctctgt  
1201 gtgtttgtgt gtatctgttg gtctccatcc tccatggggg ctacgccagg tgctgtgaca  
1261 ccccccttct gaatgaagcc ttctgacctg ggctggcact gctgggggtg aggacacatt  
1321 gccccatgag acagtcccag aacacggcag ctgctggctg tgacaatggt ttcacatcc  
1381 tttagaccaag ggatgggacc tgatgacctg ggaggactct tttagttctt acctcttgt  
1441 gttctcaata aaacagaacg

Figure 7B

9/24

## Murine LAT Nucleotide Sequence (1-1260)

1 ggcacgagca ggcggggagc aagaaagggg caggtacagc tgggcacggg gatcgtgcag  
61 ctggtagctg gggcacgggc cccagctctg gctctggggc gagcaccttt ccagagccaa  
121 cactgctctc aactcagtc agcaagagag gggagccatc cagccccgaa aggatacggc  
181 tgcctactgc cgggcggatc ccaggctgga gcccgccttg tccataccc ctgctgccac  
241 tctgtctcga ggggctgcag tgcagcaggg cctgtggcag gtgctctgca gatggaagca  
301 gacgccttga gcccgggtggg gctggggctc ctgctgctgc ccttcttggt cacgctcctg  
361 gctgccctgt gcgtgcgtg ccgtgagttg ccagtctct atgacagcac ttccacagag  
421 agtttgtacc caagaagcat cctcatcaag ccacctcaaa taaccgtccc ccgaacacct  
481 gctgtttcct accctctagt cacttcttc ccaccttga ggcagccaga cctgtctccc  
541 atcccagat ccccacagcc ccttgggggt tccatcgga tgccatctc ccagcagaat  
601 tcagatgatg ccaacagtgt ggcaagctac gagaaccagg agccagcctg taagaatgtg  
661 gatgcagatg aggatgaaga cgactatccc aacggctacc tagtggtgct gcctgacagt  
721 agtctgtctg ccgtccctgt tgtctcctct gctcctgtgc ctagcaacce tgaccttgga  
781 gacagtgcct tctctgtgga gtcgtgtgaa gattacgtga atgttctga gagtgaggag  
841 agcgcagagg cgtctctgga tgggagccgg gagtatgtga atgttctccc agagcagcag  
901 ccagtacca gggctgagct ggctctgtg aactcccagg aggtggaaga cgaaggagaa  
961 gaggaagggg tggatggaga ggaagctccc gactatgaga atctacagga gcttaactga  
1021 aagcctactg cagctgtctg tctgaaact ggacttgctg ggggtgtcgt aagaggatcc  
1081 catttgatct ctgccctgcc acagcctgag aatcttccc taacttattg tcacttggg  
1141 gtccagictg tgteccaat attctgtacc ttctgataaa gcctgagaat gaatctggtt  
1201 ccagccagac catgtcatgg aataaaggcc atgtgacata aaaaaaaaaa aaaaaaaaaa

Figure 7C

10/24

[illegible]

Human (1-233) and Murine LAT Amino Acid Sequence (1-242)

## Figure 7D

11/24

1 ggaataggtt agtttcagac aagcctgctt gccggagctc agcagacacc aggccttccg  
 61 ggcaggcctg gcccaccgtg ggccctcagag ctgctgctgg ggcattcaga accggctctc  
 121 cattggcatt gggaccagag accccgcaag tggcctgttt gcctggacat ccacctgtac  
 181 gtccccaggt ttcgggaggc ccaggggcca tgccagacct gaagctggcg ggcattggcg  
 241 tctacggcag catctcgcgt gccgaggccg aggagcacct gaagctggcg ggcattggcg  
 301 acgggctctt cctgctgcgc cagtgcctgc gctcgtggg cggtatgtg ctgtcgtcg  
 361 tgcacgatgt gcgttccac cactttcca tcgagcgcca gctcaacggc acctacgcca  
 421 ttccggcg caaagcgac ttggaccgg cagagctctg cgagtctac tcgcgcgacc  
 481 ccgacgggt gccctgcaac ctgcgcaagc cgtgcaaccg gccgtcgggc ctgagccgc  
 541 agccgggggt cttcgactgc ctgcgagacg ccatggtgcg tgactacgtg cgccagacgt  
 601 ggaagctgga gggcgaggcc ctggagcagg ccatcatcag ccaggccccg caggtggaga  
 661 agctcattgc tacgacggcc cacgagcgga tgccttgga ccacagcagc ctgacgcgtg  
 721 aggaggccga gcgcaactt tactctgggg cgcagaccga cggcaagttc ctgctgaggc  
 781 cgcggaagga gcagggcaca tacgccctgt cctcatcta tgggaagac gtgtaccact  
 841 acctcatcag ccaagacaag gcgggcaagt actgcattcc cgagggcacc aagtttgaca  
 901 cgctctggca gctggtggag tatcgaagc tgaaggcgga cgggctcctc tactgcctga  
 961 aggaggcctg ccccaacagc agtgccagca acgctcagg ggctgctgct cccacactcc  
 1021 cagcccaccc atccacgttg actcatctc agagacgaat cgacaccctc aactcagatg  
 1081 gatacacccc tgagccagca cgcataact cccagacaa accgcgccg atgccatgg  
 1141 acacgagcgt gtatgagagc cctacagcg acccagagga gctcaaggac aagaagctc  
 1201 tcctgaagcg cgataacctc ctcatagctg acattgaact tggctgcggc aactttggct  
 1261 cagtgcgcca gggcgtgtac cgcattgcga agaagcagat cgacgtggcc atcaaggtgc  
 1321 tgaagcaggg cacggagaag gcagacacgg agagatgat gcgcgaggcg cagatcatgc  
 1381 accagctgga caaccctac atcgtcggc tcattggcgt ctgccaggcc gaggcctca  
 1441 tgctggtcat ggagatggct gggggcgggc cgtgcacaa gttcctggtc ggcaagaggg  
 1501 aggagatccc tgtgagcaat gtggccgagc tgcgcacca ggtgtccatg gggatgaagt  
 1561 acctggagga gaagaactt gtgcaccgtg acctggcgcc ccgcaacgtc ctgctggtta  
 1621 accggcacta cgccaagatc agcgacttg gcctctcaa agcactgggt gccgacgaca  
 1681 gctactacac tgcccgtca gcagggaagt ggccgctcaa gtggtacgca cccgaatgca  
 1741 tcaacttccg caagtctcc agccgcagcg atgtctggag ctatggggtc accatgtggg  
 1801 aggccttgtc ctacggccag aagccctaca agaagatgaa agggccggag gtcatggcct  
 1861 tcacgagca gggcaagcgg atggagtgcc caccagagtg tccaccgaa ctgtacgac  
 1921 tcattgagtga ctgctggatc tacaagtggg aggatcgccc cgacttctg accgtggagc  
 1981 agcgcattgc agcctgttac tacagcctgg ccagcaaggt ggaagggccc ccaggcagca  
 2041 cacagaaggc tgaggctgcc tctgcctgag ctcccgtgc ccaggggagc cctccacgcc  
 2101 ggcttctccc caccctcagc cccacccag gtctgcagt ctggctgagc cctgcttgg  
 2161 tgtctccaca cacagctggg ctgtggtagg ggtgtctca ggccacaccg gccttgcat  
 2221 gcctgcctgg cccctgtcc tctctgctg gggagcaggg aggtccggga gggctcggct  
 2281 gtgcagcctg tcttgggctg gtggctcccg gagggccctg agctgagggc attgcttaca  
 2341 cggatgcctt cccctgggcc ctgacattgg agcctgggca tctcaggtg gtcaggcgta  
 2401 gatcaccaga ataaaccag ctccctctt gaaaaaaaaa aaaaaaaaaa aacc

Human ZAP-70 Nucleotide Sequence (1-2454)

Figure 8A



12/24

1 mpdpaahlpf fygsisraea eehklagma dglflrqcl rslggyvls vddvrfhhfp  
61 ierqlngtya iaggkahcgp aelcqfysqd pdglpcnlm acnrppglep qpgvfdclrd  
121 amvrdivrqt wklegdaleq aiiisqapqve kliattaher mpwyhssltr eeaerklysg  
181 qqtdgkflr prkeqgtyal slvygktvyh ylisqdkagk ycipegtkfd tlwqlveyk  
241 lkadgliyrl kevcpnssas aavaaptlpa hpstftqpqr rvdtnsdgy tpeparlass  
301 tdkprmpmpmd tsyespysd peelkdkklf lkrenllvad ielgcgnfgs vrqgvymrk  
361 kqidvaikvl kqgtekadkd emmreaqimh qldnpyivrl igvcqaealm lvmemagggp  
421 lhfllgkke ipvsnaell hqvamgmky l eeknfvhrdl aarnvllvr hyakisdfgl  
481 skalgaddsy ytarsagkwp lkwyapecin frkfssrsdv wsygvtnwea fsygqkpykk  
541 mkgpevldfi kqgkrmecpp ecppemyalm sdcwiykwed rpdfltveqr mrnyyyslas  
601 raegppqceq vacaacg

Human ZAP-70 Amino Acid Sequence (1-617)

**Figure 8B**

1 gaggaagagc cgcgggcccg gcggtgagg ccaccccggc ggcggctgga gagcgaggag  
61 gagcgggtgg ccccgcgctg cgcccgccct cgccacact ggcgagggtg gacacctgcg  
121 caggtgtgtg ccctccggcc cctgaagcat ggccagcagc ggcatggctg acagcgccaa  
181 ccacctgccc ttcttttcg gcaacatcac ccgggaggag gcagaagatt acctggtcca  
241 ggggggcatg agtcatgggc ttatttgcg gcgcagagc cgcaactacc tgggtggctt  
301 cgccctgtcc gtggccacg ggaggaaggc acaccactac accatcgagc gggagctgaa  
361 tggcacctac gccatcgccg gtggcaggac ccatgccagc ccgcccagc tctgccacta  
421 ccactcccag gagtctgatg gcctggtctg cctcctcaag aagccctca accggcccca  
481 aggggtgcag cccaagactg ggcccttga ggatttgaag gaaaacctca tcagggaata  
541 tgtgaagcag acatggaacc tgcagggtca ggctctggag caggccatca tcagtcagaa  
601 gcctcagctg gagaagctga tcgtaccac agcccatgaa aaaatgcctt ggttccatgg  
661 aaaaatctct cggaagaat ctgagcaaat tgcctgata ggatcaaga caaatggaaa  
721 gtctctgatc cgagccagag acaacaacgg ctctacgcc ctgtgcctgc tgcacgaagg  
781 gaaggtgctg cactatcgca tcgacaaaga caagacaggg aagctctcca tccccgaggg  
841 aaagaagttc gacacgctct ggagctagt cgagcattat tctataaag cagatggttt  
901 gttaagagtt ctactgtcc catgtcaaaa aatcggcaca cagggaatg ttaattttg  
961 aggccgtcca caactccag gtccccatcc tgcgtcctcc cctgcccag ggaaccggca  
1021 agagagtact gtgtcattca atccgtatga gccagaactt gcacctggg ctgcagacaa  
1081 agggccccag agagaagccc tacccatgga cacagagggtg tacgagagcc cctacgcgga  
1141 ccccgaggag atcaggccca aggaggttta cctggaccga aagctgtga cgctggaaga  
1201 caaagaactg ggctctggtg attttgaac tgtgaaaag ggctactacc aaatgaaaa  
1261 agttgtgaaa accgtggctg tgaaaatact gaaaacgag gccaatgacc ccgtcttaa  
1321 agatgagtta ttgacagaag caaatgtcat gcagcagctg gacaaccgt acatctgctg  
1381 gatgatcggg atatgagagg ccgagtcctg gatgctggtt atggagatgg cagaacttgg  
1441 tcccctcaat aagtatttc agcagaacag acatgtcaag gataagaaca tcatagaact  
1501 ggttcacag gtttccatgg gcatgaagta ctggaggag agcaattttg tgcacagaga  
1561 tctggctgca agaaatgtgt tgtagttac ccaacattac gccaagatca gtgatttcg  
1621 actttccaaa gcactgcgtg ctgatgaaa ctactacaag gccagaccc atggaaagt  
1681 gcctgtcaag tggtagctc cggatgcat caactactac aagttctcca gcaaaagcga  
1741 tgtctggagc ttggagtggt tgatgtggga agcattctcc tatgggcaga agccatctg  
1801 agggatgaaa ggaagtgaag tcaccgctat gtagagaaa ggagagcgga tgggggtccc  
1861 tgcagggtgt ccaagagaga tttacgatct catgaatctg tgctggacat acgatgtgga  
1921 aaacaggccc ggattcgag cagtggaaact gcggctgcgc aattactact atgacgtggt  
1981 gaactaaccg ctcccgacc tgcgggtggc tgccttgat cacaggagca atcacaggaa  
2041 aatgtatcca gaggaattga ttgtcagcca cctccctctg ccagtcggga gagccaggct  
2101 tggatggaac atgccacaa ctgtcacc aaagcctgtc ccaggactca cctccacaa  
2161 agcaaaggca gtcccgggag aaaagacgga tggcaggatc caaggggcta gctggattg  
2221 ttgttttct tgtctgtgtg atttccatc aggtatttt tacgatctgt ttccaaatcc  
2281 ctttcatgtc ttccacttc tctgggtccc ggggtgcatt tgttactcat cgggccagg  
2341 gacattgcag agtggcctag agcacttca cccaagcgg cttttccaa atgcccagg  
2401 atgccttagc atgtgaccc tgaagggaag gcaaaggcag aggaatttgg ctgcttctac  
2461 ggccatgaga ctgatccctg gccactgaaa agcttccctg acaataaaaa tgtttgagg  
2521 ctttaaaaag aaaaaaaaaa a

Human Syk Kinase Nucleotide Sequence (1-2541)

Figure 9A

I massgmadsa nhlpfffgni treeaedyiv qggmsdglyl lrqsrnylgg falsvahgrk  
61 ahhytierel ngtyaiaggr thaspdlch yhsqesdglv clkkpfnrp qgvqpktgpf  
121 edlkenlire yvkqtwnlqg qaleqaiisq kpqlekliat tahekmwfh gkisreeseq  
181 ivligsktng kflirardnn gsyalcilhe gkvlhyridk dktgklsipe gkkfdtlwql  
241 vehysykadg llrvltvpcq kigtqgnvnf ggrpqlpqsh passpaqgnr qestvsfnpy  
301 epelapwaad kgpqrealpm dtevyespya dpeerpkev yldrklitle dkelgsgnfg  
361 tvkkgyyqmk kvvktvavki lkneandpal kdellaeav mqqldnpyiv rmigiceaes  
421 wmlvmemael gpinkylqqn rhvkdknie lvhqvsimgmk yleesnfvr dlaarnvllv  
481 tqhyakisdf glskalrade nyykaqthgk wpvkwyapec inyykfssks dvwsfgvlmw  
541 eafsygqkpy rgmkgsevta mlekgermgc pagcpremyd lmlcwtydv enrpqfaave  
601 lrlmnyydv vn

Human Syk Kinase Amino Acid Sequence (1-612)

Figure 9B

1 gccagtgaaat tcggggggtc agccctctc cctccctcc cctgcttca ggctgctgag  
61 cactgagcag cgctcagaat ggaagccatc gccaaatatg acttcaaagc tactgcagac  
121 gacgagctga gcttcaaaag gggggacatc ctcaagggtt tgaacgaaga atgtgatcag  
181 aactggtaca aggcagagct taatggaaaa gacggcttca tcccaagaa ctacatagaa  
241 atgaaaccac atccgtggtt ttttgcaaa atcccagag ccaaggcaga agaaatgctt  
301 agcaaacagc ggcacgatgg ggccttctt atccgagaga gtgagagcgc tcctggggac  
361 ttctccctct ctgtcaagtt tggaaacgat gtgcagcact tcaagggtgct ccgagatgga  
421 gccgggaagt acttctctg ggtggtgaag ttcaattctt tgaatgagct ggtggattat  
481 cacagatcta catctgtctc cagaaaccag cagatattcc tgcgggacat agaacagggtg  
541 ccacagcagc cgacatacgt ccaggccctc ttgactttg atcccagga ggatggagag  
601 ctgggcttcc gccggggaga tttatccat gtcattgata actcagacc caactgggtg  
661 aaaggagctt gccacgggca gaccggcatg ttcccgca attatgtcac cccgtgaac  
721 cggaacgtct aagagtcaag aagcaattat taaagaaag tgaaaaatgt aaaacacata  
781 caaaagaatt aaaccacaa gctgcctctg acagcagcct gtgagggagt gcagaacacc  
841 tggccgggtc accctgtgac cctctcactt tgggtggaac tttaggggtt gggagggggc  
901 gtggattta aaaatgcaa aacttaccta taaattaaga agagtttta ttacaaattt  
961 tcaactgtgc tccttttcc cctccttgt cttttttc atcctttt ctctctgtc  
1021 catcagtga tgacgtttaa ggccacgtat agtcctagct gacgccaata ataaaaaaca  
1081 agaaacaaaa aaaaaaaac ccgaattca

Human Grb Nucleotide Sequence (1-1109)

Figure 10A

16/24

1 meaiakydfk ataddelsfk rgdilkvlne ecdqnwykae lngkdgfipk nyiemkphpw  
61 ffgkipraka eemlskqrhd gaflireses apgdfslsvk fgndvqhfkv lrdgagkyfl  
121 wvfkfnsln lvdyhrstsv srnqqiflrd ieqvpqqpty vqaldfdpq edgelgfrg  
181 dfihvmdnsd pnwwkgachg qtgmfpmyv tpvnrnv

Human Grb Amino Acid Sequence (1-217)

**Figure 10B**

17/24

1 ctaggctttt gcaaaaagct tcacgctgcc gcaagcactc agggcgcaag ggctgctaaa  
 61 ggaagcggaa cacgtagaaa gccagtcgcg agaaacgggtg ctgaccccg atgaatgtca  
 121 gctactgggc tatctggaca agggaaaacg caagcgcaaa gagaaagcag ttctgtgac  
 181 ttaagaacat tagaaccttc ctgtccacct gctgtgagaa gtccggcctc aagcggagcg  
 241 agctcttcca agcctttgac ctcttcgatg tgcaggattt tggcaaggct atctacacc  
 301 tgtctgctct gtcttgacc ccgatcgccc agaacagggg gatcatgccc tccccaccg  
 361 aggaggagag ttaggtgat gaagacatct acagtggcct gtccgaccag atcgacgaca  
 421 cgggtggagga ggatgaggac ctgtatgact gcgtggagaa tgaggaggcg gaaggcgacg  
 481 agatctatga ggacctcatg cgctcgagc ccgtgtccat gccgccaag atgacagagt  
 541 atgacaagcg ctgctgctgc ctgctggaga tccagcagac ggaggagaag tacactgaca  
 601 cgctgggctc catccagcag cattcttga agcccctgca acggttcctg aaacctcaag  
 661 acattgagat catctttatc aacattgagg acctgcttcg tgtcatact cacttctaa  
 721 aggagatgaa ggaagccctg ggcacccctg gcgcaccgaa tctctaccag gtcttcatca  
 781 aatacaagga gaggttctc gtctatggcc gctactgcag ccaggtggag tcagccagca  
 841 aacacctgga ccgtgtggcc gcagccggg aggacgtgca gatgaagctg gaggaatgtt  
 901 ctgagagagc caacaacggg aggttcactg cgcgacctgc tgatgtgccc tatgcagcga  
 961 gttctcaaat atcacctct tctccaggag ctggtgaaac acacgcagga ggcatggag  
 1021 caaggaaact gcggctggcc ctggatgcca tgagggacct ggctcagtgc gtgaacgagg  
 1081 tcaagcgaga caacgagaca ctgcgacaga tccaattt ccagctgtcc attgagaacc  
 1141 tggaccagtc tctggctcac tatggccggc ccaagatcga cggggaactc aagatcacct  
 1201 cgggtggaac gcgtccaag atggacaggt atgccttctt gctcgacaaa gctctactca  
 1261 tctgtaagcg caggggagac tctatgacc tcaaggactt tgtaaactc cacagcttcc  
 1321 aggttcggga tgactcttca ggagaccgag acaacaagaa gtggagccac atgttctcc  
 1381 tgatcgagga ccaagggtgc cagggtatg agctgttctt caagacaaga gaattgaaga  
 1441 agaagtggat ggagcagttt gagatggcca tctccaacat ctatccggag aatgccaccg  
 1501 ccaacgggca tgacttcag atgttctct ttgaggagac cacatctgc aaggcctgtc  
 1561 agatgtgct tagaggtacc tctatcagg gctaccgtg ccatcgggtc cgggcatctg  
 1621 cacacaagga gtgtctggg aggttccctc catgtggccg acatgggcaa gattccag  
 1681 gaactatgaa gaaggacaaa ctacatcgca gggctcagga caaaaagg aatgagctgg  
 1741 gtctgccccaa gatggagggtg ttccaggaat actacgggct tctccacc cctggagcca  
 1801 ttgaccctt tctacggctc aacctggag acattgtgga gtcacgaag gctgaggctg  
 1861 aacagaactg gtgggagggc agaaatacat ctactaatga aattggctgg ttcttcta  
 1921 acagggtgaa gccctatgtc catggccctc ctacaggacct gtctgttcat ctctggtacg  
 1981 caggcccat ggagcgggca ggggcagaga gcatcctggc caaccgctc gacgggactt  
 2041 tcttggtgcg gcagagggtg aaggatgcag cagaatttgc catcagcatt aaatataacg  
 2101 tcgaggtcaa gcacacggtt aaaatcatga cagcagaagg actgtaccgg atcacagaga  
 2161 aaaaggcttt ccgggggctt acggagctgg tggagttaa ccagcagaac tctctaaagg  
 2221 attgcttcaa gtctctggac accacctgc agtccccct caaggagcct gaaaagagaa  
 2281 ccatcagcag gccagcagtg ggaagcaca agtattttg cacagccaaa gcccgctatg  
 2341 acttctgcgc ccgtgaccgt tcagagctgt cgctcaagga ggggtgacat atcaagatcc  
 2401 ttaacaagaa gggacagcaa ggctgggtggc gaggggagat ctatggccgg gttggctggt  
 2461 tccctgcaa ctacgtggag gaagattatt ctgaatactg ctgagccctg gtgccttggc  
 2521 agagagacga gaaactccag gctctgagcc cggcgtggcg aggcagcga ccaggggctg  
 2581 tgacagctcc ggcgggtgga gactttggga tggactggag gaggccagcg tccagctggc  
 2641 ggtgctcccg ggatgtgccc tgacatggtt aatttataac acccgattt tctcttggg  
 2701 tcccccaag cagacggggg ctcaaggggg ttacatttaa taaaaggatg aagatgg

Human Vav Nucleotide Sequence (1-2757)

Figure 11A

1 mnvsywaiwt renasakrkq flclknirtf lstccekfgl krselfeafd lfdvqdfgkv  
61 iytlsalswt piaqnrngimp fpteesvgd ediysglsdq iddtveeded lydcveneea  
121 egdeiyedlm rsepvsmpk mteydkrccc lreiqqteek ytdtlgsiqq hflkplqrfi  
181 kpgdieiifi niedllrvht hflkemkeal gtpgapnlyq vfikykerfl vygrycsqve  
241 saskhldrva aaredvqmkl eecsqrang rftarpadga yaassqispp spgagethag  
301 gdgarklrla ldamrdlaqc vnevkrdnet lrqitnfqls ienldqslah ygrpkidgel  
361 kitsverrsk mdryafldk allickrrgd sydlkdfvnl hsfqvrddss gdrdnkkwsh  
421 mflledqga qgyelfftr elkkkwmeqf emaisniype natanghdfq mfsfeettsc  
481 kacqmlrgrt fyqgyrchrc rasahkeclg rvppcgrhgq dfpgtmkkdk lhrraqdkkr  
541 nelglpkmev fqeyyglppp pgaigpflrl npgdiveltk aaeqnwwweg mntstneigw  
601 fpcnrvkpyv hgppqdlsvh lwyagpmera gaesilanrs dgthflvrqrk kdaaeafai  
661 kynvevkhtv kimtaeglyr itekkafrgl telvefyqqn slkdcfksld tlqfpfkep  
721 ekrtrisrpa gstrygtak arydfcardr selslkegdi ikilnkkgqq gwwrgeiygr  
781 vgwfpanyve edyseyc

Human Vav Amino Acid Sequence (1-797)

Figure 11B

1 gaattccggg cccggatagc cggcggcggc ggcggcggcg gcggcggcgg cggccgggag  
 61 aggccctcc ttacgccct gcttcttcc ctgctcgca gtcagccga gccggcggac  
 121 ccgcttggg tccgacctg cccaggccat gccggcaac gtgaagaaga gctctggggc  
 181 cggggcggc acgggctccg ggggctcggg ttgggtggc ctgattggg tcatgaagga  
 241 cgccttcag ccgcaccacc accaccacca ccacctcagc cccaccgcc cggggacggg  
 301 ggacaagaag atgggtgaga agtgctgga gctcatggac aagggtggg ggttgtgca  
 361 gaacccaaag ctggcgctaa agaatagcc accctatctc ttgacctgc taccagatac  
 421 ctaccagcat ctccgtacta tctgtcaag atatgagggg aagatggaga cacttggaga  
 481 aatgagtat ttagggtgt ttatggagaa ttgatgaag aaaactaagc aaaccataag  
 541 cctctcaag gagggaaaag aaagaatga tggagagaat tctcagccta gccgaaacct  
 601 aaccaactg tccctcatct tcagccacat gctggcagaa ctaaaaggaa tcttccaag  
 661 tggactctt caggagaca catttcggat tactaaagca gatgctcgg aatttggag  
 721 aaaagcttt ggggaaaaa caatagtcct ttggaagagc ttgcacagg cctacatga  
 781 agtgcaccc atcagttctg ggctggaggc catggctctg aaatccacta ttgatctgac  
 841 ctgcaatgat tatattcgg ttttgaatt tgacatctt acccgactct ttacgccctg  
 901 gtcccttgg ctacggaatt ggaacagcct tctgttaact catcctggct acatggctt  
 961 ttgacgtat gacgaagtga aagctcggct ccagaaatc attcacaac ctggcagta  
 1021 tatctccgg ctgagctga ctgctcggg tcagtgggt attgggtatg ttactgctga  
 1081 tgggaacatt ctccagacaa tccctcaca taaacctctc ttcaagcac tgattgatg  
 1141 ctccaggaa ggcttctatt tgttctga tggacgaat cagaatcctg atctgactgg  
 1201 ctatgtgaa ccaactcccc aagaccatat caaagtgacc caggaacaat atgaattata  
 1261 ctgtgagatg ggctcccat tccaactatg taaaatatg gctgaaatg ataaggatg  
 1321 aaagattgag cctgtggac acctcatgtg cacatcctgt ctacatctt ggacggaatc  
 1381 agaaggtcag ggctgtcctt tctgccgatg tgaataaa ggtactgaac ccacgtggt  
 1441 agatcgtt gatctagag ggagtgagc cgtgtgagg caaggagcag agggagctcc  
 1501 ctcccaaat tatgatgatg atgatgatg acgagctgat gatactctt tcatgatga  
 1561 ggaattggct ggtgccaagg tggacggc gccttctcca ttccatgg cccacaagc  
 1621 ttccctccc ccggtgccac cagactga ccttctccg cagcagtat gtgtccctc  
 1681 aagtgtctt gcttggaa ctgttctaa ggctgtctt ggctccctc ataaagaca  
 1741 accattgcca gtacctcca cactcgaga tctccacca ccaccgctc cagaccggc  
 1801 atattctgt ggagcagaat cccgacctc aagacgccc ttgccttga caccaggcga  
 1861 ctgtccctc agagacaaac tgccccctgt cccctctagc cgccttgag actcatggc  
 1921 gccccggcca atcccaaaag taccagtatc tgccccagt tccagtatc cctggacagg  
 1981 aagagaatta accaaccggc actcactcc atttctatt cctcacaac tggagcccag  
 2041 accagatgtg cctaggtcga gaagcacgtt cagtctgat acctccatga gtatgaatg  
 2101 cagccatta gtaggccag agtgtagca cccaaaatc aaacctctc catctgcaa  
 2161 tgcatttat tcttggctg ccagacctc tctgtgcca aaactgccac ctggggagca  
 2221 atgtgagggt gaagaggaca cagagtacat gactccctc tccaggcctc tacggcctt  
 2281 ggatacatc cagagttcac gagcatgtga ttgcgaccag cagattgata gctgtacga  
 2341 tgaagcaatg tataatattc agtccaggc gccatctatc accgagagca gcaccttgg  
 2401 tgaagggaat ttggccgag cccatgcaa cactgtccc gaggagtcag aaataggga  
 2461 tgatgggtat gatgtccaa agccacctgt gccggccgtg ctggccgcc gaactctc  
 2521 agatatctt aatgccagc cctccttgg ctggtgtct ctggatggt atcctacaac  
 2581 aaatgtact gaagggtccc aagttcccga gaggcctcca aaaccattc cgcgagaat  
 2641 caactctga cgaaagctg gcagctgca gcaaggtagt ggtctgccg cctctgtc  
 2701 caccgctca cctcagctc ccagttagat cgagaacct atgagtcagg ggtactcta  
 2761 ccaggacatc cagaaagctt tggctattg ccagaacaac atcgagatgg ccaaaaacat  
 2821 cctccggaa tttgttcca tttcttcc tgccatgta gctacctagc acaccatc  
 2881 cctgtcag gtttagagga ccagttagt gggagttatt actcaagtgg cacctagaag  
 2941 ggcaggagt ccttggta ctacagtg aagcttgc ctctgttg gatatacat  
 3001 cagtgttcc aagattcaa agtggtaga tgaatgga gcagctagta tgtttatta  
 3061 tttatgggt ctgagtgca ttgaagggt

Human Cbl Nucleotide Sequence (1-3090)

Figure 12A



1 magnvkkssg agggsgsgsg aggliglmkd afqphhhhhh lshppctvd kkmvekcwkl  
61 mdkvvrhcqn pnvaknspp yildllpdy qhlrtvlsry egkmetlgen eyfrvfmnl  
121 mkktkqtisl fkegkermeye ensqprnlt klslifshml aelkgifpsg lfqgdtfrit  
181 kadaaefwrk afgektivpw ksfrqalhev hpiissgleam alkstidlte ndyisvfevd  
241 iftrlfqpw sllrnwnsla vthpgymafl tydevkarlq kfihkpgsyi frlsctrigq  
301 waigyvtadg nilqtiphmk plfqalidgf regfylfpdg mqnpdltgl ceptpqdhik  
361 vtqicaendk dvkiepcghl mctscitswq esegqgcpfc rceikgtepi vvdpdfprgs  
421 gslrqgaeg apspnydddd deraddslfm mkelagakve rpsspfsmap qaslppvppr  
481 ldllqrapv pastsvlgta skaasgslhk dkplpipptl rdlppppppd rpysvgaetr  
541 pqrtrlpctp gdcpsrdklp pvpssrpgds wlsrtipkvp vatpnpgdpw ngreltnrhs  
601 lpfslpsqme pradvprlgs tfsldtsmtm nsspvagpes ehpkikpsss anaiyslaar  
661 plpmpklppg eqgeseedte ymtptsrpvg vqkpepkrpl eatqssracd cdqqidscety  
721 eamytisqa lsvaensag egnlatahts tgpeesened dgydvkppv pavlartrls  
781 disnasssfg wlsldgdptn fnegsqvper ppkpfprin serkassyqq gggatanpva  
841 tapspqlsse ierlmsqgys yqdiqkalvi ahnniemakn ilrefvsiss pahvat

Human Cbl Amino Acid Sequence (1-896)

Figure 12B




Fig. 13

22/24




Fig. 14

23/24





**A****B****C**

FIG. 15

24/24



## SEQUENCE LISTING

<110> Samelson, Lawrence E.  
Zhang, Weiguo

<120> Compositions and Methods for Identifying and Testing  
Tyrosine Kinase Substrates and Their Agonists and  
Antagonists

<130> NIH-05033

<140> Unknown

<141> 1998-12-23

<150> 60/068,690

<151> 1997-12-23

<160> 15

<170> PatentIn Ver. 2.0

<210> 1

<211> 1059

<212> DNA

<213> Homo sapiens

<400> 1

```

gactctgccc ttgagggggc taggggtgca gccagcctgc tccgagctcc cctgcagatg 60
gaggaggcca tcctggtccc ctgctgctgc gggctcctgc tgctgccatc ctggccatgt 120
tgatggcact gtgtgtgcac tgccacagac tgccaggctc ctacgacagc acatcctcag 180
atagtttgta tccaaggggc atccagttca aacggcctca cacggttgcc ccctggccac 240
ctgcctaccc acctgtcacc tcctacccac ccctgagcca gccagacctg ctccccatcc 300
caagatcccc gcagcccctt gggggctccc accggacgcc atcttcccg ggggattctg 360
atgggtgcaa cagtgtggcg agctacgaga acgaggaacc agcctgtgag gatgcagatg 420
aggatgagga cgactatcac aaccacaggct acctgggtgg gcttcctgac agcaccgccg 480
ccactagcac tgctgccccca tcagctcctg cactcagcac ccctggcatc cgagacagtg 540
ccttctccat ggagtcctat gatgattacg tgaacgttcc ggagagcggg gagagcgcag 600
aagcgtctct ggatggcagc cgggagtatg tgaatgtgtc ccaggaactg catcctggag 660
cggctaagac tgagcctgcc gccctgagtt cccaggaggc agaggaagtg gaggaagagg 720
gggctccaga ttacgagaat ctgcaggagc tgaactgagg gcctgtggag gccagctctg 780
tcctggaacc aggcctgcct gggacggctg agctgggcag ctggaagtgg cctctggggtc 840
ctcacatggc gtccctgcct tgctccagcc tgacaacagc ctgagaaatc cccccgtaac 900
ttattatcac ttggtgggtt ggctgtgtgc cccogaacgc tctgcacctt ctgacgcagc 960
ctgagaatga cctgccctgg cccagccctt actctgtgta atagaataaa ggcctgcgtg 1020
tgtctgtgga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1059

```

<210> 2

<211> 1461

<212> DNA

<213> Homo sapiens

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 2

```

accccatctt catctggcct tgactctgcc cttgaggggc ctaggggtgc agccagcctg 60
ctccgagctc cctgcagat ggaggaggcc atcctggctc cctgcgtgct ggggctcctg 120
ctgctgcccc tcctggccat gttgatggca ctgtgtgtgc actgccacag actgccaggc 180
tcctacgaca gcacatctc agatagtttg tatccaagg gcatccagtt caaacggcct 240
cacacgggtg cccctgggcc acctgcctac ccacctgtca cctcctaccc acccctgagc 300
cagccagacc tgctccccat cccaagatcc ccgcagcccc ttggggggctc ccaccggacg 360
ccatctttcc cggcgggatt ctgatggtgc caacagtgtg gcgagctacg agaacgaggg 420
tgctgtctgg atccgaggtg atccaggttg gtggggagtc tggggctcgt cctggactag 480
gctgaccctt gtgtcgttac ccccagaacc agcctgtgag gatgcagatg aggatgagga 540
cgactatcac aaccacaggct acctgggtgg gcttcctgac agcaccgccg ccactagcac 600
tgctgcccca tcagctcctg cactcagcac ccctggcatc cgagacagtg cgttctccat 660
ggagtcctat gatgattacg tgaacgttcc ggagagcggg gagagcgcag aagcgtctct 720
ggatggcagc cgggagtatg tgaatgtgtc ccaggaactg catcctggag cggctaagac 780

```

```

tgagcctgcc gccctgagtt cccaggaggc agaggaagtg gaggaagagg gggctccaga 840
ttacgagaat ctgcaggagc tgaactgagg gcctgtggag gccgagtctg tcctggaacc 900
aggcctgcct gggacggcgt agctgggcag ctggaagtgg ctctgggggtc ctcacatggc 960
gtcctgccct tgctccagcc tgacaacagc ctgagaaatc cccccgtaac ttattatcac 1020
5 tttgggggttc ggctgtgttc ccccgaaacgc tctgcacctt ctgacgcagc ctgagaatga 1080
cctgccctgg cccagccctt actctgtgta atagaataaa ggctgtgctg tgtctgtgtt 1140
gagcgtgctg ctgtgtgtgc ctgtgtgcca gtctgagtca gagatttggg gatgtctctg 1200
tgtgtttgtg tgtatctgtg ggtctccatc ctccatgggg gctcagccag gtgctgtgac 1260
accccccttc tgaatgaagc cttctgacct gggctggcac tgctgggggt gaggacacat 1320
10 tgccccatga gacagtcca gaacacggca gctgctggct gtgacaatgg tttcaccatc 1380
cttagaccaa gggatgggac ctgatgacct gggaggactc ttttagttct tacctcttgt 1440
ggttctcaat aaaacagaac g                                     1461

```

&lt;210&gt; 3

&lt;211&gt; 1260

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;400&gt; 3

```

20 ggcacgagca ggcgggggagc aagaaagggg caggtacagc tgggcacggg gatcgtgcag 60
ctggtagctg gggcacgggc cccagctctg gctctggggc gagcaccttt ccagagccaa 120
cactgctctc aactcagtc agcaagagag gggagccatc cagccccgaa aggatacggc 180
tgcctactgc cgggcggatc ccaggctgga gcccgcttgg tcccataccc ctgctgccac 240
25 tctgtctcga ggggctgcag tgcagcaggg cctgtggcag gtgctctgca gatggaagca 300
gacgccttga gcccggtggg gctgggectc ctgctgctgc ccttcttggg cacgctcctg 360
gctgcccctgt gcgtgcgctg ccgtgagttg ccagtctcct atgacagcac ttccacagag 420
agtttgtacc caagaagcat cctcatcaag ccacctcaaa taaccgtccc cgaagcacct 480
gctgtttcct accctctagt cacttccttc ccacctctga ggcagccaga cctgctcccc 540
30 atcccgagat cccacagcc ccttgggggt tcccatcgga tgccatcttc ccagcagaat 600
tcagatgatg ccaacagtgt ggcaagctac gagaaccagg agccagcctg taagaatgtg 660
gatgcagatg aggatgaaga cgactatccc aacgctacc tagtggtgct gcctgacagt 720
agtctgctg ccgtccctgt tgtctcctct gctcctgtgc ctagcaaccc tgaccttgga 780
gacagtgcct tctctgtgga gtcgtgtgaa gattacgtga atgttctctga gagtgaggag 840
35 agcgcagagg cgtctctgga tgggagccgg gagtatgtga atgtgtcccc agagcagcag 900
ccagtgaaca gggctgagct ggccctctgt aactcccagg aggtggaaga cgaaggagaa 960
gaggaagggg tggatggaga ggaagctccc gactatgaga atctacagga gcttaactga 1020
aagcctactg cagctgtctg tccctgaaact ggacttgctg gggtgtcgct aagaggatcc 1080
catttgatct ctgccttgcc acagcctgag aatcttcccc taacttattg tcactttggg 1140
40 gtccagtcctg tgtccccaat attctgtacc ttctgataaa gcctgagaat gaatctgggt 1200
ccagccagac catgtcatgg aataaaggcc atgtgacata aaaaaaaaaa aaaaaaaaaa 1260

```

&lt;210&gt; 4

&lt;211&gt; 233

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

```

Met Glu Glu Ala Ile Leu Val Pro Cys Val Leu Gly Leu Leu Leu Leu
1           5           10           15

```

```

50 Pro Ile Leu Ala Met Leu Met Ala Leu Cys Val His Cys His Arg Leu
20           25           30

```

```

55 Pro Gly Ser Tyr Asp Ser Thr Ser Ser Asp Ser Leu Tyr Pro Arg Gly
35           40           45

```

```

Ile Gln Phe Lys Arg Pro His Thr Val Ala Pro Trp Pro Pro Ala Tyr
50           55           60

```

Pro Pro Val Thr Ser Tyr Pro Pro Leu Ser Gln Pro Asp Leu Leu Pro  
65 70 75 80

5 Ile Pro Arg Ser Pro Gln Pro Leu Gly Gly Ser His Arg Thr Pro Ser  
85 90 95

Ser Arg Arg Asp Ser Asp Gly Ala Asn Ser Val Ala Ser Tyr Glu Asn  
100 105 110

10 Glu Glu Pro Ala Cys Glu Asp Ala Asp Glu Asp Glu Asp Asp Tyr His  
115 120 125

Asn Pro Gly Tyr Leu Val Val Leu Pro Asp Ser Thr Pro Ala Thr Ser  
130 135 140

15 Thr Ala Ala Pro Ser Ala Pro Ala Leu Ser Thr Pro Gly Ile Arg Asp  
145 150 155 160

20 Ser Ala Phe Ser Met Glu Ser Ile Asp Asp Tyr Val Asn Val Pro Glu  
165 170 175

Ser Gly Glu Ser Ala Glu Ala Ser Leu Asp Gly Ser Arg Glu Tyr Val  
180 185 190

25 Asn Val Ser Gln Glu Leu His Pro Gly Ala Ala Lys Thr Glu Pro Ala  
195 200 205

Ala Leu Ser Ser Gln Glu Ala Glu Glu Val Glu Glu Glu Gly Ala Pro  
210 215 220

30 Asp Tyr Glu Asn Leu Gln Glu Leu Asn  
225 230

35 <210> 5  
<211> 242  
<212> DNA  
<213> Murinae gen. sp.

40 <400> 5  
meadalspvg lglilllpflv tllaalcvcrc relpvsydst steslyprsi likppqitvp 60  
rtpavsyplv tsfpplrqpdl lpiprspqp lggshrmppss qqnssddansv asyenqepac 120  
knvdateded dypngylvvl pdsspaavpv vssapvpsnp dlgsafsvs scedyvnvpe 180  
seesaeasld gsreyvvnvsp eqqpvtrel asvnsqevde egeeegvdge eapdyenlqe 240  
ln 242

45 <210> 6  
<211> 2450  
<212> DNA  
50 <213> Homo sapiens

<400> 6  
ggaatagggtt agtttcagac aagcctgctt gccggagctc agcagacacc aggccttccg 60  
ggcaggcctg gccaccgtg ggccctcagag ctgctgctgg ggcattcaga accggctctc 120  
55 cattggcatt gggaccagag accccgcaag tggcctgttt gcctggacat ccacctgtac 180  
gtccccaggt ttccgggaggc ccaggggcga tgccagaccc cgcggcgacac ctgcccttct 240  
tctacggcag catctcgctg gccgaggccg aggagcacct gaagctggcg ggcattggcg 300  
acgggctctt cctgctcgcg cagtgcctgc gctcgctggg cggctatgtg ctgtcgctcg 360  
tgacacgatgt ggccttccac cactttccca tcgagcgcca gctcaacggc acctacgcca 420  
60 ttgccggcgcg caaagcgcac tgtggaccgg cagagctctg cgagttctac tcgcgcgacc 480  
ccgacggggt gccctgcaac ctgcgcaagc cgtgcaaccg gccgtcgggc ctcgagccgc 540  
agcgggggtt cttcgactgc ctgcgagacg ccatggtgcg tgactacgtg cgccagacgt 600  
ggaagctgga gggcgaggcc ctggagcagg ccatcatcag ccaggccccg caggtggaga 660  
agctcattgc tacgacggcc cagcagcgga tgccctggta ccacagcagc ctgacgcgtg 720  
65 aggaggccga gcgcaaactt tactctgggg cgcagaccga cggcaagtgc ctgctgaggc 780  
cgcggaagga gcggggcacat acgccctgtc cctcatctat ggggaagacg tgtaccacta 840  
cctcatcagc caagacaagg cgggcaagta ctgcattccc gagggcacca agtttgacac 900  
gctctggcag ctggtggagt atctgaagct gaaggcgagc gggctcatct actgcctgaa 960



5 ggaggcctgc cccaacagca gtgccagcaa cgcctcaggg gctgctgctc ccacactccc 1020  
agcccaccca tccacgttga ctcatacctca gagacgaatc gacaccctca actcagatgg 1080  
atacaccttg agccagcacg cataacgtcc ccagacaaac cgcgcccgat gcccatggac 1140  
acgagcgtgt atgagagccc ctacagcgac ccagaggagc tcaaggacaa gaagctcttc 1200  
ctgaagcgcg ataacctcct catagctgac attgaacttg gctgcggaac ctttggctca 1260  
gtgcgccagg gcggtgtaccg catgcgcaag aagcagatcg acgtggccat caaggtgctg 1320  
aagcagggga cggagaaggc agacacggaa gagatgatgc gcgagggcga gatcatgcac 1380  
cagctggaca acccctacat cgtgcggctc attggcgctc gccaggccga ggccctcatg 1440  
ctgggtcatgg agatggctgg gggcgggccg ctgcacaagt tcctgggtcgg caagagggag 1500  
10 gagatccctg tgagcaatgt ggccgagctg ctgcaccagg tgtccatggg gatgaagtac 1560  
ctggaggaga agaactttgt gcacctgac ctggcgggccc gcaacgtcct gctgggttaa 1620  
ccggcactac gccaaagatca ggcacttggc ctctccaaag cactgggtgc cgacgacagc 1680  
tactacactg cccgctcagc aggggaagtgg ccgctcaagt ggtacgcacc cgaatgcac 1740  
aacttcgcga agttctccag ccgcagcgat gtctggagct atggggtcac catgtgggag 1800  
15 gccttgctct acggccagaa gccctacaag aagatgaaag ggccggaggt catggcgttc 1860  
atcgagcagg gcaagcggat ggagtgccca ccagagtgtc caccogaact gtacgcactc 1920  
atgagtgact gctggatcta caagtgggag gatcgccccg acttctctgac cgtggagcag 1980  
cgcatgcgag cctgttacta cagcctggcc agcaaggtgg aagggccccc aggcagcaca 2040  
cagaagctg aggcctgctg tgcctgagct cccgctgccc aggggagccc tccacgcgg 2100  
20 ctcttccccca cctcagccc caccacaggt cctgcagtct ggctgagccc tgcttggttg 2160  
tctccacaca cagctgggct gtggtagggg gtgtctcagg ccacaccggc cttgcattgc 2220  
ctgcctggcc cctgtctctc tctggctggg gagcagggag gtccgggagg gtgcggtctg 2280  
gcagcctgtc ctgggctggt ggctcccagc gggccctgag ctgagggcat tgcttacacg 2340  
gatgccttcc cctgggcccct gacattggga cctgggcatc ctgaggtggt caggcgtaga 2400  
25 tcaccagaat aaaccagct tccctcttga aaaaaaaaaa aaaaaaaacc 2450

&lt;210&gt; 7

&lt;211&gt; 617

&lt;212&gt; PRT

30 &lt;213&gt; Homo sapiens

&lt;400&gt; 7

Met Pro Asp Pro Ala Ala His Leu Pro Phe Phe Tyr Gly Ser Ile Ser  
1 5 10 15

35 Arg Ala Glu Ala Glu Glu His Leu Lys Leu Ala Gly Met Ala Asp Gly  
20 25 30

40 Leu Phe Leu Leu Arg Gln Cys Leu Arg Ser Leu Gly Gly Tyr Val Leu  
35 40 45

Ser Leu Val Asp Asp Val Arg Phe His His Phe Pro Ile Glu Arg Gln  
50 55 60

45 Leu Asn Gly Thr Tyr Ala Ile Ala Gly Gly Lys Ala His Cys Gly Pro  
65 70 75 80

Ala Glu Leu Cys Gln Phe Tyr Ser Gln Asp Pro Asp Gly Leu Pro Cys  
85 90 95

50 Asn Leu Arg Asn Ala Cys Asn Arg Pro Pro Gly Leu Glu Pro Gln Pro  
100 105 110

55 Gly Val Phe Asp Cys Leu Arg Asp Ala Met Val Arg Asp Tyr Val Arg  
115 120 125

Gln Thr Trp Lys Leu Glu Gly Asp Ala Leu Glu Gln Ala Ile Ile Ser  
130 135 140

60 Gln Ala Pro Gln Val Glu Lys Leu Ile Ala Thr Thr Ala His Glu Arg  
145 150 155 160

Met Pro Trp Tyr His Ser Ser Leu Thr Arg Glu Glu Ala Glu Arg Lys  
165 170 175

65 Leu Tyr Ser Gly Gln Gln Thr Asp Gly Lys Phe Leu Leu Arg Pro Arg  
180 185 190

Lys Glu Gln Gly Thr Tyr Ala Leu Ser Leu Val Tyr Gly Lys Thr Val  
 195 200 205  
 Tyr His Tyr Leu Ile Ser Gln Asp Lys Ala Gly Lys Tyr Cys Ile Pro  
 210 215 220  
 Glu Gly Thr Lys Phe Asp Thr Leu Trp Gln Leu Val Glu Tyr Leu Lys  
 225 230 235 240  
 Leu Lys Ala Asp Gly Leu Ile Tyr Arg Leu Lys Glu Val Cys Pro Asn  
 245 250 255  
 Ser Ser Ala Ser Ala Ala Val Ala Ala Pro Thr Leu Pro Ala His Pro  
 260 265 270  
 Ser Thr Phe Thr Gln Pro Gln Arg Arg Val Asp Thr Leu Asn Ser Asp  
 275 280 285  
 Gly Tyr Thr Pro Glu Pro Ala Arg Leu Ala Ser Ser Thr Asp Lys Pro  
 290 295 300  
 Arg Pro Met Pro Met Asp Thr Ser Val Tyr Glu Ser Pro Tyr Ser Asp  
 305 310 315 320  
 Pro Glu Glu Leu Lys Asp Lys Lys Leu Phe Leu Lys Arg Glu Asn Leu  
 325 330 335  
 Leu Val Ala Asp Ile Glu Leu Gly Cys Gly Asn Phe Gly Ser Val Arg  
 340 345 350  
 Gln Gly Val Tyr Arg Met Arg Lys Lys Gln Ile Asp Val Ala Ile Lys  
 355 360 365  
 Val Leu Lys Gln Gly Thr Glu Lys Ala Asp Lys Asp Glu Met Met Arg  
 370 375 380  
 Glu Ala Gln Ile Met His Gln Leu Asp Asn Pro Tyr Ile Val Arg Leu  
 385 390 395 400  
 Ile Gly Val Cys Gln Ala Glu Ala Leu Met Leu Val Met Glu Met Ala  
 405 410 415  
 Gly Gly Gly Pro Leu His Lys Phe Leu Leu Gly Lys Lys Glu Ile Pro  
 420 425 430  
 Val Ser Asn Val Ala Glu Leu Leu His Gln Val Ala Met Gly Met Lys  
 435 440 445  
 Tyr Leu Glu Glu Lys Asn Phe Val His Arg Asp Leu Ala Ala Arg Asn  
 450 455 460  
 Val Leu Leu Val Asn Arg His Tyr Ala Lys Ile Ser Asp Phe Gly Leu  
 465 470 475 480  
 Ser Lys Ala Leu Gly Ala Asp Asp Ser Tyr Tyr Thr Ala Arg Ser Ala  
 485 490 495  
 Gly Lys Trp Pro Leu Lys Trp Tyr Ala Pro Glu Cys Ile Asn Phe Arg  
 500 505 510  
 Lys Phe Ser Ser Arg Ser Asp Val Trp Ser Tyr Gly Val Thr Met Trp  
 515 520 525  
 Glu Ala Phe Ser Tyr Gly Gln Lys Pro Tyr Lys Lys Met Lys Gly Pro  
 530 535 540  
 Glu Val Leu Asp Phe Ile Lys Gln Gly Lys Arg Met Glu Cys Pro Pro  
 545 550 555 560

Glu Cys Pro Pro Glu Met Tyr Ala Leu Met Ser Asp Cys Trp Ile Tyr  
565 570 575

Lys Trp Glu Asp Arg Pro Asp Phe Leu Thr Val Glu Gln Arg Met Arg  
580 585 590

Asn Tyr Tyr Tyr Ser Leu Ala Ser Arg Ala Glu Gly Pro Pro Gln Cys  
595 600 605

1.0 Glu Gln Val Ala Glu Ala Ala Cys Gly  
610 615

<210> 8  
<211> 2541  
<212> DNA  
<213> Homo sapiens

20 gaggaagagc cgcgggcccg gcggtgagg ccaccccggc ggcggtgga gagcgaggag 60  
gagcggttg cccgcgctg cgcccgccct cgcctcacct ggcgcaggtg gacacctgcg 120  
cagggtgtgt cccctccggcc cctgaagcat ggccagcagc ggcatggctg acagcgccaa 180  
ccacctgccc ttcttttttcg gcaacatcac ccgggaggag gcagaagatt acctgggtcca 240  
ggggggcatg agtgatgggc ttattttgct gcgccagagc cgcaactacc tgggtggctt 300  
25 cgccctgtcc gtggcccacg ggaggaaggc acaccactac accatcgagc gggagctgaa 360  
tggcacctac gccatcgccg gtggcaggac ccatgccagc cccgcgcgacc tctgccacta 420  
ccactcccag gagtctgatg gcctgggtctg cctcctcaag aagcccttca accggcccca 480  
aggggtgcag cccaagactg ggcccttga ggatttgaag gaaaacctca tcagggaata 540  
tgtgaagcag acatggaacc tgcaggggtca ggctctggag caggccatca tcagtcagaa 600  
30 gcctcagctg gagaagctga tgcctaccac agcccatgaa aaaatgcctt ggttccatgg 660  
aaaaatctct cgggaagaat ctgagcaaat tgtcctgata ggatcaaaga caaatggaaa 720  
gttccctgat cgagccagag acaacaacgg ctctacgcc ctgtgctgc tgcacgagg 780  
gaaggtgctg cactatcgca tgcacaaaga caagacaggg aagctctcca tccccgagg 840  
aaagaagttc gacacgctct ggcagctagt cgagcattat tcttataaag cagatgggtt 900  
35 gttaagagtt ctactgtcc catgtcaaaa aatcggcaca cagggaatg ttaattttgg 960  
aggccgtcca caacttccag gttcccatcc tgcgtcctcc cctgcccagg ggaaccggca 1020  
agagagta ctgtcattca atccgtatga gccagaactt gcaccctggg ctgcagacaa 1080  
aggcccccag agagaagccc taccatgga cacagaggtg tacgagagcc cctacgcgga 1140  
ccccgaggag atcaggccca aggaggttta cctggaccga aagctgctga cgctggaaga 1200  
40 caaagaactg ggctctggta attttgaac tgtgaaaaag ggctactacc aatgaaaaa 1260  
agttgtgaaa accgtggctg tgaataact gaaaaacag gccaatgacc ccgctcttaa 1320  
agatgagtta ttagcagaag caaatgtcat gcagcagctg gacaacccgt acatcgtgcg 1380  
gatgatcggg atatgcgagg ccgagtcctg gatgctgggt atggagatgg cagaacttgg 1440  
tcccctcaat aagtatttgc agcagaacag acatgtcaag gataagaaca tcatagaact 1500  
45 gtttcatcag gtttccatgg gcatgaagta ctgggaggag agcaattttg tgcacagaga 1560  
tctggctgca agaaatgtgt tgctagttac ccaacattac gccaagatca gtgatttcgg 1620  
actttccaaa gcactgcgtg ctgatgaaaa ctactacaag gccagacccc atggaaagtg 1680  
gcctgtcaag lggtagctc cggaatgcat caactactac aagttctcca gcaaaagcga 1740  
tgtctggagc tttggagtgt tgatgtggga agcattctcc tatgggcaga agccatatcg 1800  
50 agggatgaaa ggaagtgaag tcaccgctat gttagagaaa ggagagcgga tggggtgccc 1860  
tgcaggtgt ccaagagaga tgtacgatct catgaatctg tgctggacat acgatgtgga 1920  
aaacaggccc ggattcgcag cagtggaaat gcggtcgcg aattactact atgacgtgg 1980  
gaactaaccg ctcccgcacc tgtcgtggc tgcctttgat cacaggagca atcacaggaa 2040  
aatgtatcca gaggaattga ttgtcagcca cctcctctg ccagtcggga gagccaggct 2100  
55 tggatggaac atgcccacaa cttgtcacc aaagcctgtc ccaggactca ccctccacaa 2160  
agcaaaggca gtcccgggag aaaagacgga tggcaggatc caaggggcta gctggatttg 2220  
tttgttttct tgtctgtgtg attttcatac aggttatttt tacgatctgt ttccaaatcc 2280  
ctttcacttc tttccacttc tctgggtccc ggggtgcatt tgttactcat cgggcccagg 2340  
gacattgcag agtggcctag agcactctca cccaagcgg ccttttccaa atgcccagg 2400  
60 atgccttagc atgtgactcc tgaagggaag gcaaaggcag aggaatttgg ctgcttctac 2460  
ggccatgaga ctgatccctg gccactgaaa agctttcctg acaataaaaa tgttttgagg 2520  
ctttaaaaaa a a 2541

<210> 9  
<211> 635  
<212> PRT  
<213> Homo sapiens

<400> 9  
Met Ala Ser Ser Gly Met Ala Asp Ser Ala Asn His Leu Pro Phe Phe  
1 5 10 15  
5 Phe Gly Asn Ile Thr Arg Glu Glu Ala Glu Asp Tyr Leu Val Gln Gly  
20 25 30  
Gly Met Ser Asp Gly Leu Tyr Leu Leu Arg Gln Ser Arg Asn Tyr Leu  
35 40 45  
10 Gly Gly Phe Ala Leu Ser Val Ala His Gly Arg Lys Ala His His Tyr  
50 55 60  
15 Thr Ile Glu Arg Glu Leu Asn Gly Thr Tyr Ala Ile Ala Gly Gly Arg  
65 70 75 80  
Thr His Ala Ser Pro Ala Asp Leu Cys His Tyr His Ser Gln Glu Ser  
85 90 95  
20 Asp Gly Leu Val Cys Leu Leu Lys Lys Pro Phe Asn Arg Pro Gln Gly  
100 105 110  
Val Gln Pro Lys Thr Gly Pro Phe Glu Asp Leu Lys Glu Asn Leu Ile  
115 120 125  
25 Arg Glu Tyr Val Lys Gln Thr Trp Asn Leu Gln Gly Gln Ala Leu Glu  
130 135 140  
Gln Ala Ile Ile Ser Gln Lys Pro Gln Leu Glu Lys Leu Ile Ala Thr  
145 150 155 160  
30 Thr Ala His Glu Lys Met Pro Trp Phe His Gly Lys Ile Ser Arg Glu  
165 170 175  
35 Glu Ser Glu Gln Ile Val Leu Ile Gly Ser Lys Thr Asn Gly Lys Phe  
180 185 190  
Leu Ile Arg Ala Arg Asp Asn Asn Gly Ser Tyr Ala Leu Cys Leu Leu  
195 200 205  
40 His Glu Gly Lys Val Leu His Tyr Arg Ile Asp Lys Asp Lys Thr Gly  
210 215 220  
Lys Leu Ser Ile Pro Glu Gly Lys Lys Phe Asp Thr Leu Trp Gln Leu  
225 230 235 240  
45 Val Glu His Tyr Ser Tyr Lys Ala Asp Gly Leu Leu Arg Val Leu Thr  
245 250 255  
50 Val Pro Cys Gln Lys Ile Gly Thr Gln Gly Asn Val Asn Phe Gly Gly  
260 265 270  
Arg Pro Gln Leu Pro Gly Ser His Pro Ala Thr Trp Ser Ala Gly Gly  
275 280 285  
55 Ile Ile Ser Arg Ile Lys Ser Tyr Ser Phe Pro Lys Pro Gly His Arg  
290 295 300  
60 Lys Ser Ser Pro Ala Gln Gly Asn Arg Gln Glu Ser Thr Val Ser Phe  
305 310 315 320  
Asn Pro Tyr Glu Pro Glu Leu Ala Pro Trp Ala Ala Asp Lys Gly Pro  
325 330 335  
65 Gln Arg Glu Ala Leu Pro Met Asp Thr Glu Val Tyr Glu Ser Pro Tyr  
340 345 350  
Ala Asp Pro Glu Glu Ile Arg Pro Lys Glu Val Tyr Leu Asp Arg Lys

	355	360	365	
	Leu Leu Thr Leu Glu Asp Lys Glu Leu Gly Ser Gly Asn Phe Gly Thr 370 375 380			
5	Val Lys Lys Gly Tyr Tyr Gln Met Lys Lys Val Val Lys Thr Val Ala 385 390 400			
10	Val Lys Ile Leu Lys Asn Glu Ala Asn Asp Pro Ala Leu Lys Asp Glu 405 410 415			
	Leu Leu Ala Glu Ala Asn Val Met Gln Gln Leu Asp Asn Pro Tyr Ile 420 425 430			
15	Val Arg Met Ile Gly Ile Cys Glu Ala Glu Ser Trp Met Leu Val Met 435 440 445			
	Glu Met Ala Glu Leu Gly Pro Leu Asn Lys Tyr Leu Gln Gln Asn Arg 450 455 460			
20	His Val Lys Asp Lys Asn Ile Ile Glu Leu Val His Gln Val Ser Met 465 470 475 480			
25	Gly Met Lys Tyr Leu Glu Glu Ser Asn Phe Val His Arg Asp Leu Ala 485 490 495			
	Ala Arg Asn Val Leu Leu Val Thr Gln His Tyr Ala Lys Ile Ser Asp 500 505 510			
30	Phe Gly Leu Ser Lys Ala Leu Arg Ala Asp Glu Asn Tyr Tyr Lys Ala 515 520 525			
	Gln Thr His Gly Lys Trp Pro Val Lys Trp Tyr Ala Pro Glu Cys Ile 530 535 540			
35	Asn Tyr Tyr Lys Phe Ser Ser Lys Ser Asp Val Trp Ser Phe Gly Val 545 550 555 560			
40	Leu Met Trp Glu Ala Phe Ser Tyr Gly Gln Lys Pro Tyr Arg Gly Met 565 570 575			
	Lys Gly Ser Glu Val Thr Ala Met Leu Glu Lys Gly Glu Arg Met Gly 580 585 590			
45	Cys Pro Ala Gly Cys Pro Arg Glu Met Tyr Asp Leu Met Asn Leu Cys 595 600 605			
	Trp Thr Tyr Asp Val Glu Asn Arg Pro Gly Phe Ala Ala Val Glu Leu 610 615 620			
50	Arg Leu Arg Asn Tyr Tyr Tyr Asp Val Val Asn 625 630 635			
55	<210> 10 <211> 1108 <212> DNA <213> Homo sapiens			
60	<400> 10 gccagtgaat tcggggggctc agccctctctc cctcccttcc ccctgcttca ggctgctgag 60 cactgagcag cgctcagaat ggaagccatc gccaaatatg acttcaaagc tactgcagac 120 gacgagctga gcttcaaaag gggggacatc ctcaagggtt tgaacgaaga atgtgatcag 180 aactggtaca aggagagct taatggaaaa gacggcttca ttcccaagaa ctacatagaa 240 atgaaaccac atccgtggtt ttttggcaaa atcccagag ccaaggcaga agaaatgctt 300 agcaaacagc ggcacgatgg ggcctttctt atccgagaga gtgagagcgc tcctgggggac 360 ttctccctct ctgtcaagtt tggaaacgat gtgcagcact tcaagggtgct ccgagatgga 420 gccgggaagt acttcctctg ggtggtgaag ttcaattctt tgaatgagct ggtggattat 480			
65				

cacagatcta catctgtctc cagaaaccag cagatatctc tgcgggacat agaacagggtg 540  
 ccacagcagc cgacatacgt ccaggccctc tttgacttga tccccaggag gatggagagc 600  
 tgggcttccg ccgggggagat tttatccatg tcatggataa ctcagacccc aactgggtgaa 660  
 5 aggagcttgc caccgggcaga ccggcatgtt tccccgcaat tatgtcacc cctggaaccg 720  
 gaacgtctaa gagtcaagaa gcaattatct aaagaaagtg aaaaatgtaa aacacataca 780  
 aaagaattaa acccacaagc tgcctctgac agcagcctgt gagggagtgc agaacacctg 840  
 gccgggtcac cctgtgaccc tctcactttg gttggaactt taggggggtg gagggggcgt 900  
 tggattttaa aatgccaaaa cttacctata aattaagaag agttttttatt acaaattttc 960  
 10 actgctgctc ctcttttccc ctcttttctc ttttttttca tccttttttc tcttctgtcc 1020  
 atcagtgcat gacgtttaag gccacgtata gtcctagctg acgccaataa taaaaaacia 1080  
 gaaacaaaaa aaaaaaaacc cgaattca 1108

<210> 11

<211> 176

<212> PRT

<213> Homo sapiens

<400> 11

Met Glu Ala Ile Ala Lys Tyr Asp Phe Lys Ala Thr Ala Asp Asp Glu  
 1 5 10 15

Leu Ser Phe Lys Arg Gly Asp Ile Leu Lys Val Leu Asn Glu Glu Cys  
 20 25 30

Asp Gln Asn Trp Tyr Lys Ala Glu Leu Asn Gly Lys Asp Gly Phe Ile  
 35 40 45

Pro Lys Asn Tyr Ile Glu Met Lys Pro His Pro Phe Gly Asn Asp Val  
 50 55 60

Gln His Phe Lys Val Leu Arg Asp Gly Ala Gly Lys Tyr Phe Leu Trp  
 65 70 75 80

Val Val Lys Phe Asn Ser Leu Asn Glu Leu Val Asp Tyr His Arg Ser  
 85 90 95

Thr Ser Val Ser Arg Asn Gln Gln Ile Phe Leu Arg Asp Ile Glu Gln  
 100 105 110

Val Pro Gln Gln Pro Thr Tyr Val Gln Ala Leu Phe Asp Phe Asp Pro  
 115 120 125

Gln Glu Asp Gly Glu Leu Gly Phe Arg Arg Gly Asp Phe Ile His Val  
 130 135 140

Met Asp Asn Ser Asp Pro Asn Trp Trp Lys Gly Ala Cys His Gly Gln  
 145 150 155 160

Thr Gly Met Phe Pro Arg Asn Tyr Val Thr Pro Val Asn Arg Asn Val  
 165 170 175

<210> 12

<211> 2758

<212> DNA

<213> Homo sapiens

<400> 12

ctaggctttt gcaaaaagct tcacgtctgc gcaagcactc agggcgcaag ggctgctaaa 60  
 ggaagcggaa cactagaaaa gccagtcgcg agaaacgggtg ctgaccccgg atgaatgtca 120  
 gctactgggc tatctggaca agggaaaacg caagcgcaaa gagaaagcag ttccctgtgc 180  
 ttaagaacat tagaaccttc ctgtccacct gctgtgagaa gttcggcctc aagcggagcg 240  
 agctcttcga agcctttgac ctcttcgatg tgcaggattt tggcaagggtc atctacaccc 300  
 tgtctgctct gtccctggacc ccgatcgccc agaacagggg gatcatgccc ttccccaccg 360  
 65 aggaggagag tgtaggtgat gaagacatct acagtggcct gtccgaccag atcgacgaca 420  
 cggtggagga ggatgaggac ctgtatgact gcgtggagaa tgaggaggcg gaaggcgacg 480  
 agatctatga ggacctcatg cgctcggagc ccgtgtccat gccgccaag atgacagagt 540  
 atgacaagcg ctgctgctgc ctgctgggaga tccagcagac ggaggagaag tacactgaca 600

5 cgctgggctc catccagcag catttcttga agcccttgca acgggttctg aaacctcaag 660  
 acattgagat catctttatc aacattgagg acctgcttcg tgttcatact cacttcttaa 720  
 aggagatgaa ggaagccctg ggcacccctg gcgcaccgaa tctctaccag gtcttcatca 780  
 aatacaagga gaggttctct gtctatggcc gctactgcag ccagggtggag tcagccagca 840  
 aacacctgga ccgtgtggcc gcagcccggg aggacgtgca gatgaagctg gaggaatggt 900  
 ctccagagagc caacaacggg aggttcactg cgcgacctgc tgatgggtgcc tatgcagcga 960  
 gttctcaaat atcacctcct tctccaggag ctgggtgaaac acacgcagga ggcgatggag 1020  
 caaggaaact gcggtctggc ctggatgcca tgagggacct ggctcagtgc gtgaacgagg 1080  
 tcaagcgaga caacgagaca ctgcgacaga tcaccaatct ccagctgtcc attgagaacc 1140  
 10 tggaccagtc tctggctcac tatggccggc ccaagatcga cggggaactc aagatcacct 1200  
 cgggtggaag cgcgtccaag atggacaggt atgcttctct gctcgacaaa gctctactca 1260  
 tctgtaagcg caggggagac tcctatgacc tcaaggactt tgtaaacctg cacagcttcc 1320  
 aggttcggga tgactcttca ggagaccgag acaacaagaa gtggagccac atgttctctc 1380  
 tgatcgagga ccaaggtgcc cagggctatg agctgttctt caagacaaga gaattgaaga 1440  
 15 agaagtgaat ggagcagttt gagatggcca tctccaacat ctatccggag aatgccaccg 1500  
 ccaacgggca tgacttccag atgttctcct ttgaggagac cacatcctgc aaggcctgtc 1560  
 agatgctgct tagaggtacc ttctatcagg gctaccgctg ccacggtgac cgggcatctg 1620  
 cacacaagga gtgtctgggg agggctccctc catgtggccg acatgggcaa gatttcccag 1680  
 gaactatgaa gaaggacaaa ctacatcgca gggctcagga caaaaagagg aatgagctgg 1740  
 20 gtctgccccaa gatggaggtg tttcaggaat actacgggct tcctccaccc cctggagcca 1800  
 ttggaccctt tctacggctc aaccctggag acattgtgga gctcacgaag gctgaggctg 1860  
 aacagaactg gtgggagggc agaaatacat ctactaatga aattggctgg tttccttgta 1920  
 acagggtgaa gccctatgtc catggccctc ctacggacct gctgtttcat ctctgtacg 1980  
 caggcccat ggagcgggca ggggcagaga gcatcctggc caaccgctcg gacgggactt 2040  
 25 tcttggtgcg gcagaggggtg aaggatgcag cagaatttgc catcagcatt aaatataacg 2100  
 tcgaggtcaa gcacacgggt aaaatcatga cagcagaagg actgtaccgg atcacagaga 2160  
 aaaaggcttt ccgggggctt acggagctgg tggagtttta ccagcagaac tctctaaagg 2220  
 attgcttcaa gtctctggac accaccttgc agttcccctt caaggagcct gaaaagagaa 2280  
 ccacagcag gccagcagtg ggaagcacia agtatttttg cacagccaaa gcccgctatg 2340  
 30 acttctgcgc ccgtgaccgt tcagagctgt cgctcaagga ggggtgacatc atcaagatcc 2400  
 ttaacaagaa gggacagcaa ggctgggtggc gaggggagat ctatggccgg gttggctggt 2460  
 tccctgccaa ctacgtggag gaagattatt ctgaatactg ctgagccctg gtgcttggc 2520  
 agagagacga gaaactccag gctctgagcc cggcgtggcg aggcagcgga ccaggggctg 2580  
 tgacagctcc ggccgggtgga gactttggga tggactggag gaggccagcg tccagctggc 2640  
 35 ggtgctcccg ggatgtgccc tgacatggtt aatttataac accccgattt tctcttggg 2700  
 tcccctcaag cagacggggg ctcaaggggg ttacatttta ataaaaggat gaagatgg 2758

&lt;210&gt; 13

&lt;211&gt; 797

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

Met Asn Val Ser Tyr Trp Ala Ile Trp Thr Arg Glu Asn Ala Ser Ala

1

5

10

15

Lys Arg Lys Gln Phe Leu Cys Leu Lys Asn Ile Arg Thr Phe Leu Ser

20

25

30

Thr Cys Cys Glu Lys Phe Gly Leu Lys Arg Ser Glu Leu Phe Glu Ala

35

40

45

Phe Asp Leu Phe Asp Val Gln Asp Phe Gly Lys Val Ile Tyr Thr Leu

50

55

60

Ser Ala Leu Ala Trp Thr Pro Ile Ala Gln Asn Arg Gly Ile Met Pro

65

70

75

80

Phe Pro Thr Glu Glu Glu Ser Val Gly Asp Glu Asp Ile Tyr Ser Gly

85

90

95

Leu Ser Asp Gln Ile Asp Asp Thr Val Glu Glu Asp Glu Asp Leu Tyr

100

105

110

Asp Cys Val Glu Asn Glu Glu Ala Glu Gly Asp Glu Ile Tyr Glu Asp

115

120

125

Leu Met Arg Ser Glu Pro Val Ser Met Pro Pro Lys Met Thr Glu Tyr

	130	135	140
	Asp Lys Arg Cys Cys Cys Leu Arg Glu Ile Gln Gln Thr Glu Glu Lys		
	145	150	155 160
5	Tyr Thr Asp Thr Leu Gly Ser Ile Gln Gln His Phe Leu Lys Pro Leu		
		165	170 175
10	Gln Arg Phe Leu Lys Pro Gln Asp Ile Glu Ile Ile Phe Ile Asn Ile		
		180	185 190
	Glu Asp Leu Leu Arg Val His Thr His Phe Leu Lys Glu Met Lys Glu		
		195	200 205
15	Ala Leu Gly Thr Pro Gly Ala Pro Asn Leu Tyr Gln Val Phe Ile Lys		
		210	215 220
	Tyr Lys Glu Arg Phe Leu Val Tyr Gly Arg Tyr Cys Ser Gln Val Glu		
		225	230 235 240
20	Ser Ala Ser Lys His Leu Asp Arg Val Ala Ala Ala Arg Glu Asp Val		
		245	250 255
25	Gln Met Lys Leu Glu Glu Cys Ser Gln Arg Ala Asn Asn Gly Arg Phe		
		260	265 270
	Thr Ala Arg Pro Ala Asp Gly Ala Tyr Ala Ala Ser Ser Gln Ile Ser		
		275	280 285
30	Pro Pro Ser Pro Gly Ala Gly Glu Thr His Ala Gly Gly Asp Gly Ala		
		290	295 300
	Arg Lys Leu Arg Leu Ala Leu Asp Ala Met Arg Asp Leu Ala Gln Cys		
		305	310 315 320
35	Val Asn Glu Val Lys Arg Asp Asn Glu Thr Leu Arg Gln Ile Thr Asn		
		325	330 335
40	Phe Gln Leu Ser Ile Glu Asn Leu Asp Gln Ser Leu Ala His Tyr Gly		
		340	345 350
	Arg Pro Lys Ile Asp Gly Glu Leu Lys Ile Thr Ser Val Glu Arg Arg		
		355	360 365
45	Ser Lys Met Asp Arg Tyr Ala Phe Leu Leu Asp Lys Ala Leu Leu Ile		
		370	375 380
	Cys Lys Arg Arg Gly Asp Ser Tyr Asp Leu Lys Asp Phe Val Asn Leu		
		385	390 395 400
50	His Ser Phe Gln Val Arg Asp Asp Ser Ser Gly Asp Arg Asp Asn Lys		
		405	410 415
55	Lys Trp Ser His Met Phe Leu Leu Ile Glu Asp Gln Gly Ala Gln Gly		
		420	425 430
	Tyr Glu Leu Phe Phe Lys Thr Arg Glu Leu Lys Lys Lys Trp Met Glu		
		435	440 445
60	Gln Phe Glu Met Ala Ile Ser Asn Ile Tyr Pro Glu Asn Ala Thr Ala		
		450	455 460
	Asn Gly His Asp Phe Gln Met Phe Ser Phe Glu Glu Thr Thr Ser Cys		
		465	470 475 480
65	Lys Ala Cys Gln Met Leu Leu Arg Gly Thr Phe Tyr Gln Gly Tyr Arg		
		485	490 495



Cys His Arg Cys Arg Ala Ser Ala His Lys Glu Cys Leu Gly Arg Val  
 500 505 510  
 Pro Pro Cys Gly Arg His Gly Gln Asp Phe Pro Gly Thr Met Lys Lys  
 515 520 525  
 Asp Lys Leu His Arg Arg Ala Gln Asp Lys Lys Arg Asn Glu Leu Gly  
 530 535 540  
 Leu Pro Lys Met Glu Val Phe Gln Glu Tyr Tyr Gly Leu Pro Pro Pro  
 545 550 555 560  
 Pro Gly Ala Ile Gly Pro Phe Leu Arg Leu Asn Pro Gly Asp Ile Val  
 565 570 575  
 Glu Leu Thr Lys Ala Glu Ala Glu Gln Asn Trp Trp Glu Gly Arg Asn  
 580 585 590  
 Thr Ser Thr Asn Glu Ile Gly Trp Phe Pro Cys Asn Arg Val Lys Pro  
 595 600 605  
 Tyr Val His Gly Pro Pro Gln Asp Leu Ser Val His Leu Trp Tyr Ala  
 610 615 620  
 Gly Pro Met Glu Arg Ala Gly Ala Glu Ser Ile Leu Ala Asn Arg Ser  
 625 630 635 640  
 Asp Gly Thr Phe Leu Val Arg Gln Arg Val Lys Asp Ala Ala Glu Phe  
 645 650 655  
 Ala Ile Ser Ile Lys Tyr Asn Val Glu Val Lys His Thr Val Lys Ile  
 660 665 670  
 Met Thr Ala Glu Gly Leu Tyr Arg Ile Thr Glu Lys Lys Ala Phe Arg  
 675 680 685  
 Gly Leu Thr Glu Leu Val Glu Phe Tyr Gln Gln Asn Ser Leu Lys Asp  
 690 695 700  
 Cys Phe Lys Ser Leu Asp Thr Thr Leu Gln Phe Pro Phe Lys Glu Pro  
 705 710 715 720  
 Glu Lys Arg Thr Ile Ser Arg Pro Ala Val Gly Ser Thr Lys Tyr Phe  
 725 730 735  
 Gly Thr Ala Lys Ala Arg Tyr Asp Phe Cys Ala Arg Asp Arg Ser Glu  
 740 745 750  
 Leu Ser Leu Lys Glu Gly Asp Ile Ile Lys Ile Leu Asn Lys Lys Gly  
 755 760 765  
 Gln Gln Gly Trp Trp Arg Gly Glu Ile Tyr Gly Arg Val Gly Trp Phe  
 770 775 780  
 Pro Ala Asn Tyr Val Glu Glu Asp Tyr Ser Glu Tyr Cys  
 785 790 795  
 <210> 14  
 <211> 3090  
 <212> DNA  
 <213> Homo sapiens  
 <400> 14  
 gaattccggg cccggatagc cggcggcggc ggcggcggcg gcggcggcgg cggccgggag 60  
 aggccctcc ttcacgcct gcttctctcc ctgcctcgca gtcgagccga gccggcgagac 120  
 ccgcctgggc tccgacctg cccaggccat ggccggcaac gtgaagaaga gctctggggc 180  
 cgggggcggc acgggctccg ggggctcggg ttcgggtggc ctgattgggc tcatgaagga 240

cgccttccag cgcaccacc accaccacca ccacctcagc ccccaaccgc cggggacggt 300  
 ggacaagaag atggtggaga agtgctggaa gctcatggac aaggtggtgc ggttgtgtca 360  
 gaacccaaag ctggcgctaa agaatagccc accttatatc ttagacctgc taccagatac 420  
 ctaccagcat ctccgtacta tcttgtcaag atatgagggg aagatggaga cacttggaga 480  
 5 aaatgagtat tttaggggtgt ttatggagaa tttgatgaag aaaactaagc aaaccataag 540  
 cctcttcaag gagggaaaag aaagaatgta tgaggagaat tctcagccta ggcgaaacct 600  
 aaccaactgt tccctcatct tcagccacat gctggcagaa ctaaaaggaa tctttccaag 660  
 tggactcttt cagggagaca catttcggat tactaaagca gatgctgcgg aattttggag 720  
 10 aaaagctttt ggggaaaaga caatagtccc ttggaagagc tttcgacagg ctctacatga 780  
 agtgcatccc atcagttctg ggctggagge catggctctg aaatccacta ttgatctgac 840  
 ctgcaatgat tatatttcgg tttttgaatt tgacatcttt acccgactct ttcagcctg 900  
 gtctctcttt ctcaggaatt ggaacagcct tgctgtaact catcctggct acatggcttt 960  
 tttgacgtat gacgaagtga aagctcggct ccagaaattc attcacaac ctggcagtta 1020  
 tatcttccgg ctgagctgta ctgctctggg tcagtgggct attgggtatg ttactgctga 1080  
 15 tgggaacatt tccagacaa tccctcaca caacctctc ttccaagcac tgattgatgg 1140  
 cttcagggaa ggcttctatt tgtttcctga tggacgaaat cagaatcctg atctgactgg 1200  
 cttatgtgaa ccaactcccc aagaccatat caaagtgacc caggaacaat atgaattata 1260  
 ctgtgagatg ggctccacat tccaactatg taaaatatgt gctgaaaatg ataaggatgt 1320  
 aaagattgag ccctgtggac acctcatgtg cactcatcct ctacatcct ggcaggatc 1380  
 20 agaaggtcag ggctgtcctt tctgcccag tgaaattaaa ggtactgaac ccatcgtgg 1440  
 agatccgttt gatcctagag ggagtggcag cctgttgagg caaggagcag agggagctcc 1500  
 ctccccaat tatgatgatg atgatgatga acgagctgat gatactctct tcatgatga 1560  
 ggaattggct ggtgccaagg tggaaacggc gccttctcca ttctccatgg cccacaagc 1620  
 25 ttcccttccc cgggtgccac cagacttga cctctgccc cagcagatgt gtgttccctc 1680  
 aagtgtctct gctcttggaa ctgcttctaa ggctgtctct ggctcccttc ataaagacaa 1740  
 accattgcca gtacctccca cacttcgaga tcttccacca ccaccgcctc cagaccggcc 1800  
 atattctgtt ggagcagaat cccgacctca aagacgcccc ttgcttctga caccaggcga 1860  
 ctgtccctcc agagacaaac tgcccctgt cccctctagc cgccttggag actcatggct 1920  
 gcccggcca atcccaaaag taccagtatc tgccccaagt tccagtgatc cctggacagg 1980  
 30 aagagaatta accaaccggc actcacttcc attttcatgt ccctcacaaa tggagcccag 2040  
 accagatgtg cctaggctcg gaagcacgtt cagtctggat acctccatga gtatgaatag 2100  
 cagcccatga gtaggtccag agtgtagca ccccaaaatc aaaccttct catctgcca 2160  
 tgccatttat tctctggctg ccagacctct tctgtgcca aaactgccac ctggggagca 2220  
 atgtgagggg gaagaggaca cagagtacat gactccctct tccaggcctc tacggccttt 2280  
 35 ggatacatcc cagagttcac gagcatgtga ttgcgaccag cagattgata gctgtacgta 2340  
 tgaagcaatg tataatatc agtccaggc gccatctatc accgagagca gcacctttg 2400  
 tgaagggaa ttggccgca cccatgccaa cactggctcc gaggagtcag aaaatgagga 2460  
 tgatgggtat gatgtccaa agccacctgt gccggccgtg ctggcccgc gaactctctc 2520  
 agatatctct aatgccagct cctcctttgg ctggttgtct ctggatgggt atectacaac 2580  
 40 aaatgtcact gaaggttccc aagttcccga gaggcctcca aaaccattcc cgcggagaat 2640  
 caactctgaa cggaaagctg gcagctgtca gcaaggtagt ggtcctgccc cctctgctgc 2700  
 caccgcctca cctcagctct ccagtgaat cgagaacctc atgagtcagg ggtactccta 2760  
 ccaggacatc cagaaagctt tggctcattgc ccagaacaac atcgagatgg ccaaaaacat 2820  
 cctccgggaa tttgtttcca tttcttctcc tgcccattga gctacctagc acaccatctc 2880  
 45 cctgctgcag gtttagagga ccagtqagtt gggagttatt actcaagtgg cacctagaag 2940  
 ggcaggagtt cctttggtga cttcacagtg aagtcttgcc ctctctgtgg gatatcacat 3000  
 cagtggttcc aagatttcaa agtggtgaaa tgaaaatgga gcagctagta tgttttatta 3060  
 ttttatgggt cttgagtga tttgaaggtg 3090

50 <210> 15  
 <211> 906  
 <212> PRT  
 <213> Homo sapiens

55 \* <400> 15  
 Met Ala Gly Asn Val Lys Lys Ser Ser Gly Ala Gly Gly Gly Thr Gly  
 1 5 10 15

60 \* Ser Gly Gly Ser Gly Ser Gly Gly Leu Ile Gly Leu Met Lys Asp Ala  
 20 25 30

BNSDOCID: &lt;WO\_\_\_9932627A2\_1\_&gt;

Cys Thr Ser Cys Leu Thr Ser Trp Gln Glu Ser Glu Gly Gln Gly Cys  
 405 410 415  
 5 Pro Phe Cys Arg Cys Glu Ile Lys Gly Thr Glu Pro Ile Val Val Asp  
 420 425 430  
 Pro Phe Asp Pro Arg Gly Ser Gly Ser Leu Leu Arg Gln Gly Ala Glu  
 435 440 445  
 10 Gly Ala Pro Ser Pro Asn Tyr Asp Asp Asp Asp Glu Arg Ala Asp  
 450 455 460  
 Asp Thr Leu Phe Met Met Lys Glu Leu Ala Gly Ala Lys Val Glu Arg  
 465 470 475 480  
 15 Pro Pro Ser Pro Phe Ser Met Ala Pro Gln Ala Ser Leu Pro Pro Val  
 485 490 495  
 Pro Pro Arg Leu Asp Leu Leu Pro Gln Arg Val Cys Val Pro Ser Ser  
 500 505 510  
 20 Ala Ser Ala Leu Gly Thr Ala Ser Lys Ala Ala Ser Gly Ser Leu His  
 515 520 525  
 25 Lys Asp Lys Pro Leu Pro Val Pro Pro Thr Leu Arg Asp Leu Pro Pro  
 530 535 540  
 Pro Pro Pro Pro Asp Arg Pro Tyr Ser Val Gly Ala Glu Ser Arg Pro  
 545 550 555 560  
 30 Gln Arg Arg Pro Leu Pro Cys Thr Pro Gly Asp Cys Pro Ser Arg Asp  
 565 570 575  
 35 Lys Leu Pro Pro Val Pro Ser Ser Arg Leu Gly Asp Ser Trp Leu Pro  
 580 585 590  
 Arg Pro Ile Pro Lys Val Pro Val Ser Ala Pro Ser Ser Ser Asp Pro  
 595 600 605  
 40 Trp Thr Gly Arg Glu Leu Thr Asn Arg His Ser Leu Pro Phe Ser Leu  
 610 615 620  
 Pro Ser Gln Met Glu Pro Arg Pro Asp Val Pro Arg Leu Gly Ser Thr  
 625 630 635 640  
 45 Phe Ser Leu Asp Thr Ser Met Ser Met Asn Ser Ser Pro Leu Val Gly  
 645 650 655  
 Pro Glu Cys Asp His Pro Lys Ile Lys Pro Ser Ser Ser Ala Asn Ala  
 660 665 670  
 50 Ile Tyr Ser Leu Ala Ala Arg Pro Leu Pro Val Pro Lys Leu Pro Pro  
 675 680 685  
 55 Gly Glu Gln Cys Glu Gly Glu Glu Asp Thr Glu Tyr Met Thr Pro Ser  
 690 695 700  
 Ser Arg Pro Leu Arg Pro Leu Asp Thr Ser Gln Ser Ser Arg Ala Cys  
 705 710 715 720  
 60 Asp Cys Asp Gln Gln Ile Asp Ser Cys Thr Tyr Glu Ala Met Tyr Asn  
 725 730 735  
 65 Ile Gln Ser Gln Ala Pro Ser Ile Thr Glu Ser Ser Thr Phe Gly Glu  
 740 745 750  
 Gly Asn Leu Ala Ala Ala His Ala Asn Thr Gly Pro Glu Glu Ser Glu  
 755 760 765

Asn Glu Asp Asp Gly Tyr Asp Val Pro Lys Pro Pro Val Pro Ala Val  
770 775 780

5 Leu Ala Arg Arg Thr Leu Ser Asp Ile Ser Asn Ala Ser Ser Ser Phe  
785 790 795 800

Gly Trp Leu Ser Leu Asp Gly Asp Pro Thr Thr Asn Val Thr Glu Gly  
805 810 815

10 Ser Gln Val Pro Glu Arg Pro Pro Lys Pro Phe Pro Arg Arg Ile Asn  
820 825 830

Ser Glu Arg Lys Ala Gly Ser Cys Gln Gln Gly Ser Gly Pro Ala Ala  
835 840 845

15 Ser Ala Ala Thr Ala Ser Pro Gln Leu Ser Ser Glu Ile Glu Asn Leu  
850 855 860

20 Met Ser Gln Gly Tyr Ser Tyr Gln Asp Ile Gln Lys Ala Leu Val Ile  
865 870 875 880

Ala Gln Asn Asn Ile Glu Met Ala Lys Asn Ile Leu Arg Glu Phe Val  
885 890 895

25 Ser Ile Ser Ser Pro Ala His Val Ala Thr  
900 905



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, C07K 14/705, 16/28, G01N 33/50, C12N 15/62, G01N 33/53, C12Q 1/68</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/32627</b>
		<b>(43) International Publication Date:</b> 1 July 1999 (01.07.99)
<b>(21) International Application Number:</b> PCT/US98/27400	<b>(74) Agents:</b> CARROLL, Peter, G. et al.; Medlen & Carroll, LLP, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US).	
<b>(22) International Filing Date:</b> 23 December 1998 (23.12.98)		
<b>(30) Priority Data:</b> 60/068,690 23 December 1997 (23.12.97) US		
<b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 60/068,690 (CIP) Filed on 23 December 1997 (23.12.97)	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
<b>(71) Applicant (for all designated States except US):</b> THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US).	<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SAMELSON, Lawrence, E. [US/US]; 6707 East Avenue, Chevy Chase, MD 20815 (US). ZHANG, Weiguo [-/US]; 259 Congressional Lane #410, Rockville, MD 20852 (US).	<b>(88) Date of publication of the international search report:</b> 25 November 1999 (25.11.99)	

(54) Title: THE PROTEIN TYROSINE KINASE SUBSTRATE LAT AND ITS USE IN THE IDENTIFICATION OF (ANT)AGONISTS OF THE KINASE

```

human LAT      1  MEPAILVPCVLGLLLLPIL.AMLMALCVHCHRLPQS.DSTSSDSLPRGI 49
                ||..| |..||||||:|.:| |||:|: ||..| ||||:| ||:|
murine LAT     1  MEADALSPVGLGLLLLPFLVTLAALCVRCRLPVS.DSTSTESLPRSI 50
                ||..| |..||||||:|.:| |||:|: ||..| ||||:| ||:|
50  QKRPHTVAPWFFA..|PVVTSYFPPLSQPDLLPIPRSPQPLQGSHRTFSS 97
                :|.|:|...:|.|| | |||:|:| | | | | | | | | | | | | |
51  LIKFPQITVPRTAVE|PLVTSFPPLRQPDLLPIPRSPQPLQGSHRTFSS 100
                :|.|:|...:|.|| | |||:|:| | | | | | | | | | | | | |
98  RRDSDGANSVAS|ENKEPACK..DADEDEDD|HNFG|LVVLDPDSTPATST 145
                ..:|:|:|:|:|:|:| | | | | | | | | | | | | | | | | |
101 QQNSDDANSVAS|ENQEPACKNVDADEDEDD|PN.G|LVVLDPDSSPAAVP 149
                ..:|:|:|:|:|:|:| | | | | | | | | | | | | | | | | |
146 AAPSPALSTPGIRDSAFSMESIDD|VNVPESEGESAEASLDGSR|VNVV 195
                ...|||. |.:| | |||:|:|:| | | | | | | | | | | | | |
150 VVSSAPVPSNPDGDSAFSVESCED|VNVPESEGESAEASLDGSR|VNVV 199
                ...|||. |.:| | |||:|:|:| | | | | | | | | | | | | |
196 QELHPGAAKTEPAALSSQEAE.....VEEGAPD|ENLQELN 233
                .|:| | | | | | | | | | | | | | | | | | | | | | |
200 PEQOP.VTRARLASVNSQVEDEGEEREGVQGEAPD|ENLQELN 242
                .|:| | | | | | | | | | | | | | | | | | | | | | |

```

**Human (1-233) and Murine LAT Amino Acid Sequence (1-242)**

**(57) Abstract**

The invention generally relates to compositions and methods for identifying and testing tyrosine kinase signaling pathway agonists and antagonists, and more particularly, methods and compositions for screening compounds and identifying compounds that will modulate the interaction of protein tyrosine kinase substrates with their intracellular ligands, as well as between their intracellular ligands and other members of the signaling pathway.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/27400

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 C07K16/28 G01N33/50 C12N15/62  
G01N33/53 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SIEH, M. ET AL.: "Grb2 and phospholipase C-gammal associate with a 36- to 38-kilodalton phosphotyrosine protein after T-cell receptor stimulation." MOLECULAR AND CELLULAR BIOLOGY, vol. 14, no. 7, July 1994 (1994-07), pages 4435-42, XP002117612 the whole document	1-12
X	--- DATABASE EMBL - EMBEST21 'Online! Entry HS1325208, Acc.no.AA535611, 26 July 1997 (1997-07-26) STRAUSBERG, R. ET AL.: "nf87d04.s1 NCI_CGAP_Co3 Homo sapiens cDNA clone IMAGE:926887." XP002117615 the whole document --- -/--	11



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

5 October 1999

Date of mailing of the international search report

18/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/27400

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	ZHANG, W. ET AL.: "LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation." CELL, vol. 92, 9 January 1998 (1998-01-09), pages 83-92, XP002117613 the whole document	1-25
P,X	WEBER, J.R. ET AL.: "Molecular cloning of the cDNA encoding pp36, a tyrosine-phosphorylated adapter protein selectively expressed by T cells and natural killer cells." JOURNAL OF EXPERIMENTAL MEDICIN, vol. 187, 18 April 1998 (1998-04-18), pages 1157-61, XP002117614 the whole document	1-13,17, 18,20-25

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/27400

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
See FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98 /27400

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

The antibody of claim 4 could be an antibody against a portion of a fusion protein, which comprises a seq.ID.4, but whereby the antibody is directed against a portion, not derived from seq.ID.4. A meaningful search for the whole of claim 4 was therefore not possible. Consequently, the search has been limited to antibodies against proteins comprising a substantial portion of seq.ID.4.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.